

Phytochemical screening and antibacterial activity of medicinal plants used to treat typhoid fever in Bamboutos division, West Cameroon

Tsobou Roger¹, Mapongmetsem Pierre-Marie², Voukeng Kenfack Igor³, Van Damme Patrick⁴

¹Department of Plant Biology, University of Dschang, P.O.Box 67 Dschang, Cameroon; Department of Biological Sciences, University of Ngaoundéré, P.O.Box 454 Ngaoundéré, Cameroon. ²Department of Plant Production, Laboratory of Tropical and Subtropical Agriculture and Ethnobotany, ³Department of Biological Sciences, University of Ngaoundéré, P.O.Box 454 Ngaoundéré, Cameroon. ⁴Department of Biochemistry, Faculty of Science, University of Dschang; Dschang, Cameroon.

ARTICLE INFO

Article history:

Received on: 02/03/2015

Revised on: 13/04/2015

Accepted on: 30/04/2015

Available online: 27/06/2015

Key words:

Phytochemical screening, antimicrobial, typhoid fever, medicinal plants, MIC, bactericidal.

ABSTRACT

This study was undertaken to document how typhoid is traditionally treated in Bamboutos division. For this purpose thirty eight plants species were selected. These plants underwent phytochemical screening and antibacterial study using standard procedures. The antibacterial tests using agar well diffusion method and microdilution assay indicated that, all the thirty eight plant samples showed activity against *S. typhi*, while *S. paratyphi A* and *S. paratyphi B* reacted on fifteen and fourteen plants respectively. The highest zones of inhibition were obtained from *Senna alata* with diameter of 24, 22.5 and 20.5 mm against *S. paratyphi A*, *S. paratyphi B* and *S. typhi* respectively at 160 mg/ml concentration. The lowest MIC values 128 µg/ml was exhibited by the extract of *Vitex doniana* against *Salmonella paratyphi A*. Bactericidal activity was obtained by the extract of *Carica papaya*, *Pseudarthria confertiflora*, *Moringa oleifera* and *Harungana madagascariensis*. Antibacterial screened of *Pseudarthria confertiflora* was reported for the first time. *Annona muricata*, *Lagdera alata*, *Spathodea campanulata*, *Cordia platythyrsa*, *Carica papaya*, *Terminalia glaucescens* and *Pseudarthria confertiflora* gave positive results for all secondary metabolites while other plants contained two to five metabolites. The presence of these secondary metabolites probably contributes to the antibacterial potential of these plants. This finding supported the uses of these plants for treatment of typhoid fever and other infectious diseases in the study area.

INTRODUCTION

Knowledge on plant uses is the result of many years of man's interaction and selection on the most desirable, the most vigorous and the most successful plant present in the immediate environment at a given time (Rindos, 1984). According to the World Health Organization, 80% of people in developing countries still depend on local medicinal plants to fulfill their primary health needs (WHO, 2002). Besides that, there is a global consensus on the benefits of phytopharmacy and at present medicinal plants occupy a key position in plant research and medicine. In many African countries, such as Ghana, Mali, Nigeria and Zambia, the first line of treatment for 60% of the

children with high fevers, is the use of herbal medicines at home (WHO, 2003). The importance of plants in medicine remains even of greater relevance with the current global shift to obtain drugs from plants sources, as a result of which attention has been given to the medicinal value of herbal remedies for safety, efficacy and economy (Abubakar *et al.* 2009).

Plants constitute an important source of active ingredients which differ widely in terms of structure and therapeutic properties. The continued investigation into the secondary plant metabolites for anti-infective properties has gained importance in recent years because of the alarming increase in resistance of pathogenic microorganisms to existing antibiotic. For instance, the emergence and spread of *Salmonella* resistance to many commonly used antibiotics (Ciprofloxacin, Ampicillin, Chloromphenicol, Amoxicillin) is now a subject of international concern.

* Corresponding Author

Tsobou Roger, University of Dschang, Faculty of Science,
P O BOX. 67; Phone: 6 75 72 30 72. Email: tsobou Roger@yahoo.fr

The problem has become endemic in many developing countries, causing enormous childhood morbidity and high cost of treatment (Leume, 1999). Multidrug resistant *Salmonella* species are being increasingly reported from the developed world. There is therefore, the need for efficient and safe vaccine which can be used as a preventive public health tool (Leume, 1999).

Thus, the resistances of *Salmonella* to these antibiotics couple with the high cost of treatment have prompted the present study on plant used for the treatment of typhoid fever. Furthermore, typhoid fever is a systemic infection caused by the bacterium *Salmonella enterica* subspecies *enterica* serotype *typhi*, which is acquired by ingestion of contaminated food and water (Iroha *et al.*, 2010).

Much more, in 2009, Zofou *et al.*, report that this disease remains the second cause of mortality in the Bamboutos Division after malaria. In the same way, the report provided by the NGO "Action for the formation and the natural stock management", indicated that, 2715 cases of typhoid fever and 1998 cases of diarrheas and gastro-enteritis were recorded in 2010 in this Division. In the same line, Nanfack *et al.*, in 2014 have found that typhoid fever was the hydrous widespread disease in this locality, follow by dysentery and gastro-enteritis. Besides, each year the disease affects at least 12.6 million people world-wide, most of them resides in the developing countries of Southeast Asia and Africa (WHO, 2003).

Typhoid fever is uncommon in industrialized regions such as the USA, Canada, Europe, Australia and Japan and new cases of the disease in these countries are related to a journey in the developing countries (Papadimitropoulos *et al.*, 2004). Mortality rates associated with this disease vary from one region to other, with highest reported from Indonesia, Nigeria and India (Miller *et al.*, 1994).

The prevalence of this disease and the increasing prices of medicine have resulted in the demand for discovery of less expensive but more potent sources of drugs. Plants are one of the best sources of potent drugs. In fact, medicinal plants represent a rich source from which antimicrobial agents may be obtained (Kubmarawa *et al.*, 2007).

The traditional knowledge on medicinal plants that is inherent within local communities is a very important source of information that continually provides the present-day herbal remedies (Shahid-ud-Daula & Basher, 2009). In recent years, researches in various countries have been directed towards discovering the medicinal uses of plants through phytochemical and antimicrobial screening (Hashim *et al.*, 2010).

Nevertheless, the present investigation represents a preliminary screening of some medicinal plants used against typhoid fever in Bamboutos division.

These medicinal plants have been claimed by traditional medical practitioners in this locality to be effective when used for the treatment of typhoid fever. However, no detailed reports on anti-typhoid activity of these plants exist in literature in Cameroon; therefore, the present work investigates phytochemical compounds and antibacterial activity against *Salmonella* species.

MATERIALS AND METHODS

Study area

The Bamboutos Division is located in the western highlands and extends between 5–6° N and between 9–11°E (Figure 1). It's one of the eight divisions which forms the West Region of Cameroon and is bordered to the north by Mezam Division, to the south by Mifi and Menoua divisions, and to the west by Noun Division. The Bamboutos division covers an area of 1155 km² which represents 8.31% of the total area of the West Region of Cameroon. Located on the eastern slope of the Bambouto Mountain from which it is named, Bamboutos Division is characterized by the great diversity of its relief, climate, vegetation, and soils. Mount Bambouto is the third highest mountain in Cameroon (2740 m) after Mount Cameroon (Fako) (4100 m) and Mount Oku (3008 m). The main occupation in the community is farming. There are two seasons: the dry season from November to March and the rainy season from March to October. The climate is subtropical with an annual rainfall estimated at 1621.5 mm and a mean annual temperature of 24–29°C (MINEF, 1999). They have four main languages used in the community (Ngomba'a, *Ngiembon*, *Ngombalé* and *Megaka*).

Collection of ethnomedical information

An ethnobotanical survey was carried out in the Bamboutos Division January–November 2009. Traditional healers and elderly persons were the target key informants in the study, and the selection process was based on the knowledge base of informants, experience, and current practices in ethnobotany medicine of the target individual. The traditional healers were interviewed using semi-structured questionnaires as described by Martin (1995) and open-ended conversations. Trips were made to the sites where traditional healers usually go to harvest plants. The interviews and discussions were carried out in the local language for each of the villages visited. Since the author is a native of the division, data on the local names of the plants, the plant parts used, mode of usage and administration, and mode of preparation were easily obtained. Selected respondents were well-known in the community due to their long practice in providing services related to traditional health care. The age of the respondents ranged between 40 and 95 years (29 were between 40-50, whereas 18 were between 51-60 and 23 were above 60 years old). Forty of them were traditional healers, whereas the rest were elders who had gained knowledge on medicinal uses of plants from their parents and relatives who were historically using the plants.

Collection of plant samples

Fresh plant (s) parts that were found to be used by traditional healers and elderly people for treatment of typhoid fever after carrying out the survey by use of questionnaires were collected from Bamboutos division as shown in Figure 1. The taxonomic identities of these plants were confirmed by the senior taxonomist at the Yaoundé National Herbarium and the voucher specimens were kept in the Department of Plant Biology, University of Dschang.

(Washington, 1995). The wells were then filled with 50 μ l of each ethanolic extract at a concentration of 160 mg/ml, 80 mg/ml and 40 mg/ml. Also, concentration of 10 μ g/ml of ciprofloxacin and 200 μ g/ml DMSO were introduced into other wells as positive and negative controls, respectively with the aid of micropipette. The plates were incubated at 37°C for 24 h. antibacterial activity was determined by measuring the zone of inhibition surrounding the well. The zones of inhibition were then measured, recorded and compared with standard control, Ciprofloxacin (10 μ g/ml). The assays were carried out under aseptic conditions. All tests were performed in triplicate.

Determination of MIC and MBC

Preparation of the solutions of extracts

The stock ethanolic extracts solutions were prepared by dissolving 59 mg of each extracts in 200 μ l of 10% dimethylsulphoxide (DMSO) and mixed with 1600 μ l of Mueller Hinton broth, to give the final concentration of 8192 μ g/ml and serially diluted two fold to obtain the concentration ranges from 128-8192 μ g/ml for the extract and 2-256 μ g/ml for ciprofloxacin. 100 μ l of each concentration was added in 96 wells plates containing 100 μ l of MHB and inoculums standardized at 1.5×10^8 CFU/ml. Negative growth controls were included in every test. The plates were covered with a sterile plate sealer, then agitated to mix the contents of the wells using a plate shaker and incubated at 37°C for 24 h. The MIC of each sample was detected following addition (40 μ l) of 2% *p*-iodonitrotetrazolium chloride and incubation at 37°C for 30 minutes (Pettit *et al.*, 2005). Viable microorganisms reduced the yellow dye to a pink colour. MIC was defined as the lowest sample concentration that prevented this change and exhibited complete inhibition of bacterial growth. For the determination of MBC, a portion of liquid (50 μ l) from each well that showed no change in color was placed on MHB (150 μ l each new 96 wells plates) and incubated at 37°C for 24 h. The lowest concentration that yielded no growth after this sub-culturing was taken as the MBC. The MBC/MIC was calculated to determine if the extract is bactericidal or bacteriostatic, according to Noumedem *et al.*, (2013).

STATISTICAL ANALYSIS

Data were expressed as mean \pm standard deviation. The data obtained were subjected to ANOVA test to determine whether there was significant difference in zone of inhibition between extract concentrations and also antibiotic used.

RESULTS

Seventy informants (traditional healers and elderly persons) were interviewed on how they used plants to treat typhoid fever. Fifty-nine species, distributed among 56 genera and 33 families were recorded (Tsobou *et al.*, 2013). Thirty-eight of the 59 species recorded were selected for this study (Table 1). Various parts were harvested depending on the parts the informants used in

the treatment of typhoid fever. The leaves, barks, whole plant, roots, fruits and stem were the ones that were harvested, but the part that is used most was found to be the leaves, followed by stem bark. The most common mode of preparation was decoction. The species name, family, vernacular name, ethnomedical uses elsewhere, plant part used, biological activity and chemical studies of the 38 selected species are presented in Table 1. However, *Bidens pilosa* was the most cited plant in our study. The only mode of administration was oral.

Phytochemical screening

Phytochemical screening of collected plant species has been carried out following the methods reported in literature and the results have been reported in table 2. The result of the phytochemical analysis showed that, these plants extracts are rich in at least one of the following class of compounds: triterpenes, flavonoids, alkaloids, polyphenols, sterols and saponins. All these group of compounds were found in *Annona muricata*, *Picalima nitida*, *Laggera alata*, *Spathodea campanulata*, *Cordia platythyrsa*, *Carica papaya*, *Terminalia glaucescens*, *Pseudarthria confertiflora*, *Senna alata* and *Musa paradisiaca*. Triterpenes were found in all plant extracts except *Gossypium barbadense* and *Dioscorea dumetorium*. While alkaloids have been found to be absent in 18 species (i.e. *Pseudospondias microcarpa*, *Stereospermum acuminatissimum*, *Cupressus lusitanica*, *Entada abyssinica*, *Theobroma cacao* and *Mitracarpus villosus*). Polyphenols have been found to be present in thirty seven species. Flavonoids are generally present in most of the 38 plants except in *Dracaena deisteliana*, *Pteridium aquilinum*, *Dioscorea dumetorium*, *Ocimum gratissimum* and *Paullinia pinnata*. Sterols and saponins are present in 36 and 23 plants species respectively.

Antibacterial activity

The results of the antibacterial screening of the ethanolic extract on the test strains are shown in table 2. All the thirty eight (38) plants tested showed activity against *S. typhi*. While 23 and 24 plant species did not showed any activity against *S. paratyphi A* and *S. paratyphi B* respectively. The zone of inhibition increased with increasing concentration of extract in wells. This showed that the concentration influence the activity. The highest zone of inhibition were obtained from *Senna alata* against *Salmonella paratyphi A*, *S. paratyphi B* and *S. typhi* with diameter of 24, 22.5 and 20.5 mm respectively at 160 mg/ml.

Among 38 plants tested *Pseudarthria confertiflora*, *Terminalia glaucescens*, *Senna alata*, *Dacryodes edulis* and *Stereospermum acuminatissimum* were found to be the most effective against *S. typhi*. *Senna alata* and *Rauvolfia vomitoria* were found to be the most sensitive against *Salmonella paratyphi A*. Similarly, *Senna alata*, *Pseudarthria confertiflora* and *Rauvolfia vomitoria* were found to be the most active against *S. paratyphi B*. The inhibition zone (7.5 – 24 mm) of each extract on all the bacterial used was less than the zone of inhibition caused by ciprofloxacin (Table 2).

Table 1: Information about medicinal plants recorded and tested in this study.

Species	Freq	Local names	Reported relevant ethnomedical uses elsewhere	Part used/MP	Biological activity	Chemical constituents
<i>Eremomastax speciosa</i> (Hochst) Cufod Acanthaceae	20	Kouokmegar, pankuzem, panzemmock, panzemmock, piezeumok	Dysentery, anemia, irregular menstruation, fractures cough, constipation, pile and urinary infections (Adjanohoun <i>et al.</i> , 1996). Infertility, burns, post-partum and fungi infections (Oben <i>et al.</i> , 2006).	L (I, D)	Antidiarrheic properties (Oben <i>et al.</i> , 2006).	Tannins, phenols, flavonoids, alkaloids, saponins, terpenes (Mboso <i>et al.</i> , 2013).
<i>Pseudospondias microcarpa</i> Engl. Anacardiaceae	2	Gueme	Teeth problems (Adjanohoun <i>et al.</i> , 1996).	B (D)		Alkaloids, tannins, terpenoids, steroids and cardiotoniques heterosides
<i>Annona muricata</i> Linn. Annonoaceae	5	Corossole, saba saba sour-sop	Hernia, intestinal worms, breast lactation (Baskar <i>et al.</i> , 2006). Hypertension and stomach pains (Anbu <i>et al.</i> , 2010).	L, B (D)	Antidiarrheic activity, antispasmodic, sedatif, antitumor and antibacterial activity (Baskar <i>et al.</i> , 2006). Cytotoxic, molluscidic, anti-inflammatory and anthelmintic (Pimenta <i>et al.</i> , 2011).	Alkaloids, flavonoids, glycosides, reductive sugar, steroids and tannins
<i>Picralima nitida</i> (Stapf) Th. et H. Apocynaceae	10	Djicka	Malaria, male impotence, dysmenorrhoea, digestive tract infections (Adjanohoun <i>et al.</i> , 1996). Inflammations, pneumonia and intestinal worms (Ubulom <i>et al.</i> , 2011).	R, F (D, M)	Antimicrobial, antifungal, antiprotozoa (Ubulom <i>et al.</i> , 2011).	Alkaloids, cardiotonic glycosides, saponins, tannins and terpens (Ubulom <i>et al.</i> , 2011)
<i>Rauvolfia vomitoria</i> Afz. Apocynaceae	5	Not found	Snake bites, madness, malaria (Bellomaria & Kacou, 1995). Hypertension (N'Guessan <i>et al.</i> , 2009). Jaundice, gonorrhoea, schizophrenia, intestinal worms and urinary infections (Adjanohoun <i>et al.</i> , 1996).	B, R (D)	Anticancer activity, antipyretic effect, antihypertensives, sedatives, emetic, purgative, dysenteric, arbotive and insecticidal properties (Oyewole & Massaquoi, 2008).	Active indole alkaloids, β -carboline alkaloids and alstonine (Oyewole & Massaquoi, 2008).
<i>Dracaena deisteliana</i> Engl Asparagaceae	15	Kikeng, kion	Cough (Adjanohoun <i>et al.</i> , 1996)	L, R (D, I, S)	Antileishmaniales, antimalaria, antimolluscidic, fungistatic and fungicidic (Okunji <i>et al.</i> , 1996).	Not found
<i>Bidens pilosa</i> Linn. Asteraceae	40	Lietmik, lipiliep metsemik	Cough, malaria, expulsion of the placenta, fractures, panaris and convulsions (Adjanohoun <i>et al.</i> , 1996). Diabetes, dysentery diarrhoea (Rabe & Van Staden, 1997). Diabetes, liver diseases, malaria (Brandao <i>et al.</i> , 1997).	Wp, L (D, I)	Anti-hyperglycemiales (Ubilas <i>et al.</i> , 2000). Anti-ulcerogenic (Tan <i>et al.</i> , 2000). Anti-inflammatory (Jager <i>et al.</i> , 1996). Hypotensives, anti-pyretic, anticancerous and anti-tumor, antivirales (Chiang <i>et al.</i> , 2007). Antifungal (Deba <i>et al.</i> , 2008). Antibacterial (Rojas <i>et al.</i> , 2006).	Flavonoids, phenylacetylenes, alkaloids, steroids, triterpenoids and tannins (Khan <i>et al.</i> , 2001). Saturated carbohydrate, aliphatic carboxylic acids, acetylenic 38 hydrocarbons, phenols, chalcones, flavonols, porphyrines (Silva <i>et al.</i> , 2011)
<i>Emilia coccinea</i> (Sims) G. Don. Asteraceae	18	Made-sonlume	Male and female infertility (preorgasm ejaculation, pelvic inflammatory disease), gastric and dysmenorrhoea (Focho <i>et al.</i> , 2009). Cough (Ubulom, 2010). Fever (Burkill, 1985).	Wp (D, C)	Antimicrobial activities of <i>E. sonchifolia</i> (Yoga Latha <i>et al.</i> , 2009).	Alkaloids, tannins, steroids, terpenoids, flavonoids and cardiac glycosids (Edeoga <i>et al.</i> , 2005).
<i>Laggera alata</i> (G. Don) Sch. Bip Ex. Oliv. Asteraceae	7	Depack-kenan, negikock	Hepatitis, arthritis, bronchitis, nephritis (Wu <i>et al.</i> , 2011). Fever (Adjanohoun <i>et al.</i> , 1996).	L (D)	Antibacterial, anti-inflammatory and anti-leukaemia (Asfaw <i>et al.</i> , 2003).	Essential oils (Asfaw <i>et al.</i> , 2003); eremophiloids eudesmanoids (Zhao <i>et al.</i> , 2003). Thymoquinol dimethyl ether & α -eudesmol (Ekundayo <i>et al.</i> , 1989). Flavonoids (Wu <i>et al.</i> , 2006).
<i>Vernonia colorata</i> (Willd.) Drake Asteraceae	2	Bitali, mekang, melute	Diabetes, toxoplasmosis, worms infections (Benoit <i>et al.</i> , 2000). Fever, infertility, epilepsy, rheumatism, wounds, gonorrhoea, hepatitis & digestive tract infections (Arbonnier, 2002).	L (D, S)	Antimicrobial (Kelmanson <i>et al.</i> , 2000). Antimalarial, antispasmodic (Benoit <i>et al.</i> , 1996).	Oils & sesquiterpens (Kerharo & Adam, 1974). Vernolide & hydrovernalide (Gasquet & Bamba, 1985).

<i>Spathodea campanulata</i> P.Beauv. Bignoniaceae	3	Foufougue, foukfouk	Hernia, schizophrenia, leucorrhoea, malaria & haemorrhoids (Adjanohoun <i>et al.</i> , 1996). Infertility, postpartum & weakness impotence (Focho <i>et al.</i> , 2009). Dysentery, ulcer, gonorrhoea, diarrhoea & poisons (Amusan <i>et al.</i> , 1995). Diabetes (Nagla'a, 2007). Haemorrhoids (Agyare <i>et al.</i> , 2009).	L, B (D)	Antimalarial, hypoglycemic, anti-HIV (Niyonzima <i>et al.</i> , 1999). Antimicrobial, antioxidant & nephroprotective (Rajesh <i>et al.</i> , 2010).	Spathodic acids, steroids, saponins, ursolic acids, tomentosolic acids, verminosids, pectic compounds, anthocyanins (Amusan <i>et al.</i> , 1996). Phenolic acids, caffeic acids, ferulic acids, flavonoids (Nagla'a, 2007). Specioside, diglucoside kampeferol & caffeic acids (Elusiyan <i>et al.</i> , 2011).
<i>Stereospermum acuminatissimum</i> K. Schum. Bignoniaceae	2	Watefè	Not found	B (D, M)	Not found	Not found
<i>Cordia platythyrsa</i> Bak. Boraginaceae	4	Fapbè	Not found	F (S)	Not found	Not found
<i>Canarium Schweinfurthii</i> Engl. Burseraceae	3	Beré, pui	Cough (Adjanohoun <i>et al.</i> , 1996). Fever, constipation, malaria, diarrhoea, sexual infections, rheumatism (Koudou <i>et al.</i> , 2005).	B (D)	Antioxydant, antimicrobial (Obame <i>et al.</i> , 2008).	Lipids and fatty acids (Obame <i>et al.</i> , 2008).
<i>Dacryodes edulis</i> (G.Don) H.J. Burseraceae	8	Zo'o	Digestive tract infections, ear infections, dysentery & anemia (Ayuk <i>et al.</i> , 1999). Leprosy (Bouquet, 1969). Skin infections (Ajibesin <i>et al.</i> , 2008).	L, B (D)	Antimalarial (Zofou <i>et al.</i> , 2011). Antibacterial (Ajibesin <i>et al.</i> , 2011). Antimicrobial & antioxydant (Obame <i>et al.</i> , 2011).	Essential oils rich in monoterpenes, sesquiterpenes, diterpenes, triterpenes, phellandrène & β -caryophyllène (Onocha <i>et al.</i> , 1999).
<i>Carica papaya</i> Linn. Caricaceae	27	Papaye, popo'o	Asthma, rhinitis (Morton, 1977). Wounds, burns, dyspepsia, gastro-intestinales disorders, cancer, infectious diseases (Otsuki <i>et al.</i> , 2010). Diabetes, malaria, gonorrhoea, jaundice, headache, splenomegaly (Adjanohoun <i>et al.</i> , 1996). Urinary tract infections (Mesfin <i>et al.</i> , 2005). Hypertension, galactagogue, dyspepsia (Arbonnier, 2002).	L, B, F (D)	Anthelmintic, antimicrobial, anti-tumor, antiprotozoa (Banerjee, 2001). Antioxydant, anti-inflammatory (Imaga <i>et al.</i> , 2010).	Folic acids, vitamin B12, alkaloids, saponins, glycosides, tannins, anthraquinones, flavonoids (Suresh <i>et al.</i> , 2008). Carpaine, carpasemine, chymopapaine (Gurub-Fakim, 2008).
<i>Terminalia glaucescens</i> Planch. Combretaceae	11	Tsi-sa'a	Diabetes, bacteria infections, obesity, typhoid fever, malaria, asthma, hepatosplenomegaly (Adjanohoun <i>et al.</i> , 1996). Dysentery, anti-HIV (Koudou <i>et al.</i> , 1995).	B (D)	Antimicrobial (Ndukwe <i>et al.</i> , 2005; Balou <i>et al.</i> , 2010). Antispasmodic, antivirale, antioxydants (Njomen <i>et al.</i> , 2011).	Glaucinoic acids, triterpenoids (Atta-Ur-Rahmana <i>et al.</i> , 2005). Alkaloids, saponins, tannins, steroids, flavonoids, anthraquinones, phlobatannins (Adebayo & Ishola, 2009; Kolapo <i>et al.</i> , 2009).
<i>Cupressus lusitanica</i> Mill. Cupressaceae	5	Sapin	Postpartum, hair loss (Bussmann & Ashley, 2010).	Wp, L, S (D)	Not found	Not found
<i>Pteridium aquilinum</i> (Linn.) kulh. Dennstaedtiaceae		Koukoumazong	Antiprotozoa (Crane, 1990).	L (D)	Antimicrobial activity (Hassan <i>et al.</i> , 2007).	Glycoside anthraquinones, cardiac glycosides, cyanogenic glycosides, essential oils (Hassan <i>et al.</i> , 2007).
<i>Dioscorea dumentorium</i> Dioscoreaceae	3	Leliock, meliock neliock	Not found	L (D)	Not found	Not found
<i>Entada abyssinica</i> Rich. Fabaceae	A. 3	Not found	Infertility, menstruation disorders (Chifundera, 1998). Asthma, tuberculosis, cough (Otieno <i>et al.</i> , 2011). Rheumatism, abdominale pains, diarrhoea, fever (Bekele-Tesemma <i>et al.</i> , 1993). Malaria, syphilis, wounds, psoriasis, convulsions (Arbonnier, 2002).	B (D)	Antivirale (Cos <i>et al.</i> , 2002). Antibacterial (Fabry <i>et al.</i> , 1998). Antiprotozoa (Freiburghaus <i>et al.</i> , 1998).	5S, 6R, 8a R-5-(carboxymethyl)-3-4-4a, 5, 6, 7, 8, 8a-Octahydro-5, 6, 8a trimethylnaphthalenecarboxylic, methyl 3,4,5-trihydroxybenzoate, 1,2,3-triol benzene, 2,3-dihydroxypropyltriacetate (Teke <i>et al.</i> , 2011).

<i>Pseudarthria confertiflora</i> (A. Rich.) Baker Fabaceae	2	Not found	Not found	L (D)	Not found	Not found
<i>Senna alata</i> (Linn.) Roxb Fabaceae	21	Foupan	Constipation, gastro-enteritis, intestinal worms, eczema, typhoid fever, hepatitis, jaundice (Adjanohoun, <i>et al.</i> , 1996). Skin infections, syphilis, gonorrhoea (Arbonnier, 2002). Hypertension (Okafor & Ham, 1999).	L (D)	Hepatoprotective (Pieme <i>et al.</i> , 2006). Antimicrobial (Awal <i>et al.</i> , 2004).	Phenols, tannins, saponins, alkaloids, steroids, flavonoids, carbohydrates (Owoyale <i>et al.</i> , 2005).
<i>Harungana madagascariensis</i> Lam-expoir Hypericaceae	5	Not found	Menstruation disorders, dysentery, female infertility, hernia (Adjanohoun <i>et al.</i> , 2009). Postpartum (Focho <i>et al.</i> , 2009). Malaria, fever, nervous depression, kidney & liver diseases (Adeneye <i>et al.</i> , 2008).	B (D)	Antimicrobial activity (Okoli <i>et al.</i> , 2002). Analgesic, anti-inflammatory, antioxydants (Kouam <i>et al.</i> , 2006). Antiplasmodic (Ndjakou Lenta <i>et al.</i> , 2007).	flavonoids, alkaloids, saponins, glycosides, tannins (Moulari <i>et al.</i> 2006). Harunmadagascarins C & D, Kenganthrol D, harumadagascarins A & B, harunganol B, harungin anthrone, bazouanthrone (Kouam <i>et al.</i> , 2007).
<i>Ocimum gratissimum</i> Linn. Lamiaceae	15	Kotemadjo, masep, masepo	Gonorrhoea, vaginitis, metritis (Abdulrahma, 1992). Respiratory infections, diarrhoea, headache, conjunctivitis, skin infections, pneumonia, fever (Okigbo & Ogbonnanya, 2006). Cough, convulsion, gastroenteritis (Adjanohoun <i>et al.</i> , 1996). Hypertension, malaria (N'Guessan <i>et al.</i> , 2009). Facilitate the childbirth (Kamatenesi <i>et al.</i> , 2007).	L (D)	Antimicrobial, antidiabetic activity (Oussou <i>et al.</i> , 2008). Insecticidal activity (Seri-Kouassi <i>et al.</i> , 2004). Anti-HIV activity (Elujoba, 2000).	Sterols, polyterpens, polyphenols, flavonoids, catechic tannins, alkaloids (N'Guessan <i>et al.</i> , 2009). Monoterpens, sesquiterpens, saponosids (Oussou <i>et al.</i> , 2008).
<i>Vitex doniana</i> Sweet Lamiaceae	6	Vounetane, voutane	Typhoid fever (Faleyimi <i>et al.</i> , 2012). Syphilis (Adjanohoun <i>et al.</i> , 1996). Dysmenorrhoea, infertility, gastritis, smallpox, epilepsy, wounds, snake bites (Arbonnier, 2002).	B (D)	Antimicrobial activity (Kubmarawa <i>et al.</i> , 2007; Iroha <i>et al.</i> , 2010).	Sponins, tannins, essential oils (Kubmarawa <i>et al.</i> , 2007).
<i>Gossypium barbadense</i> Linn. Malvaceae	10	Coton, ti-sisi	Pelvic pains, postpartum (Rajith <i>et al.</i> , 2010). Malaria (Ajaiyeoba <i>et al.</i> , 2006).	L (D)	Not found	Not found
<i>Theobroma cacao</i> L. Malvaceae	2	Caca	Wounds (Agyare <i>et al.</i> , 2009)	L, B (D)	Not found	Not found
<i>Thespesia populnea</i> (Linn.) soland ex corr Malvaceae	6	Kepfou	Asthma, cholera, diarrhoea, diabetes, dysentery, gonorrhoea, haemorrhoids, hernia, wounds (Parekh & Chanda, 2008). Liver diseases (Venkata & Sai, 2011).	L (S)	Antidiabetic, antioxydant activity (Parthasarathy <i>et al.</i> , 2009). Antimicrobial activity (Saravanakumar <i>et al.</i> , 2009).	Alkaloids, flavonoids, carbohydrates, phytosterols, tannins, saponins, proteins, amino acids, terpens, phenols, mucilage (Venkata et Sai, 2011). Essential oils, glycosides, fatty acids (Siddharth <i>et al.</i> , 2009). Gossypol, herbacetine, kaempferol (Parekh & Chanda, 2008). Thespones, thespesones, mansonone-D & H, gossypol (Akila & Rani, 1993).
<i>Ficus sur</i> Forssk Moraceae	5	Gack, gaia	Galactagogue (Addo-Fordjou <i>et al.</i> , 2008). Wounds, burns, ulcers (Agyare <i>et al.</i> , 2009).	B (D)	Not found	Not found
<i>Moringa oleifera</i> Lam. Moringaceae	2	Moringa	Typhoid fever (Faleyimi <i>et al.</i> , 2012). Skin infections, headache, rheumatism (Amri & Kisangau, 2012). Urinary tract infections, diarrhoea, dysentery, diabete, hypertension (Fahey, 2005).	L (D)	Antidiabetic, hypocholesterolemic activity, anti-inflammatory, liver, kidney infections, cardiovascular problems, anticancer (Bharali <i>et al.</i> , 2003).	Gallic tannins, catechol tannins, steroids, triterpenoids, flavonoids, saponins, anthraquinones, alkaloids (Kasolo <i>et al.</i> , 2010).

<i>Musa paradisiaca</i> Linn. Musaceae	22	Kedong, kadong	Diarrhoea, dysentery, intestinal infections, diabetes, uraemia, nephritis, hypertension, venereal diseases (Ghani, 2003). Nervous depression, epilepsy, hysteria, mastitis, cystitis, prostatitis, syphilis (Focho <i>et al.</i> , 2009).	L (D)	Antimicrobial, hematoprotectives, hypoglycemic, antihypertensive activity (Biswas <i>et al.</i> , 2011).	Pectines, flavonoids (Ragasa <i>et al.</i> , 2007). Serotonine, norepinephrine, tryptophane, indolic compounds, tannins, starch, iron, cristalized & non cristalized sugar, vitamin C, vitamin B, albuminoids, fatty acids, mineral salts (Ghani, 2003). Alkaloids, steroids, flavonoids, saponins, sugars, tannins (Biswas <i>et al.</i> , 2011).
<i>Psidium guajava</i> Linn. Myrtaceae	22	Goyave, gravou	Weakness impotence (Focho <i>et al.</i> , 2009). Diarrhoea, dysentery, dermatitis (Adjanooun <i>et al.</i> , 1996). Vaginitis, haemorrhoids, leucorrhoea (Arbonnier, 2002). Malaria, infectious diseases (Gurib-Fakim, 2008).	L, B (D)	Antimicrobial activity (Gurib-Fakim, 2008). Antioxydants (Ayoola <i>et al.</i> , 2008). Hepatoprotective, antispasmodic, hypoglycemiatic, hypotensive (Ojewole, 2005).	Tannins, essential oils, cinnamic acids, 3-hexenoic, quercetin, glycosides, eugenol (Gurib-Fakim, 2008).
<i>Imperata cylindrica</i> (L.)Rausch. Poaceae	4	Kenick, neuck, nick	Wounds (Anbu <i>et al.</i> , 2009). Abdominale pains (Kamatenesi <i>et al.</i> , 2011).	Wp (D)	Antibacterial, antipyretic (Balangcod <i>et al.</i> , 2012).	Alkaloids, flavonoids, tannins, glycosides (Balangcod <i>et al.</i> , 2012).
<i>Paullinia pinnata</i> Linn. Sapindaceae	3	Dzick	Rheumatism, weakness impotence (Addo-Fordjour <i>et al.</i> , 2008). Ulcers, haemorrhoids, wounds (Agyare <i>et al.</i> , 2009). Malaria (Adinorthey <i>et al.</i> , 2012). Fever, hypertension, contraceptive (N'Guessan <i>et al.</i> , 2009). Buruli ulcer (Yemoa <i>et al.</i> , 2008). Syphilis (Focho <i>et al.</i> , 2009).	L (D)	Stimulent, antioxydants (Annan & Houghton, 2010).	Tannins, saponins (Yemoa <i>et al.</i> , 2008). Sterols, polyterpens, polyphenols, flavonoids, tannins, alkaloids, saponosids (N'Guessan <i>et al.</i> , 2009).
<i>Gardenia ternifolia</i> Schum et Thonn. Rubiaceae	2	Metoucbouor	Ascites, diarrhoea, rachitism, leprosy, constipation, hepatitis, rheumatism, onchocerciasis, female infertility, wounds, haemorrhoids (Arbonnier, 2002).	B (D)	Purgatives, cholagogue, diuretic, emetic, asthenia (Arbonnier, 2002).	Not found
<i>Mitracarpus villosus</i> (Swartz)DC. Rubiaceae	3	Not found	Leprosy, skin infections (Jegade <i>et al.</i> , 2005). Dysentery, diarrhoea (Dalziel, 1937).	Wp (D)	Antibacterial, anti-inflammatory (Jegade <i>et al.</i> , 2005).	Stigmasterols, urloic acids (Sofowora, 1986). Coumarins (Ekpendu, 1995).
<i>Lantana camara</i> Linn. Verbenaceae	7	Lantana	Cataracte, snake bites, epilepsy (Kamatenesi <i>et al.</i> , 2011). Malaria (Clarkson <i>et al.</i> , 2004). diarrhoea, cough, asthma, jaundice, constipation, gastritis, asthenia, rheumatism, arthritis (Arbonnier, 2002).	L (D)	Antimicrobial (Tripathi, 2010). Antiulcerogenic, antihyperglycemic, antimotility, anti-inflammatory, antifertility, anticancer, antiproliferative (Ghosh & Sarma, 2010). Hemolitic activity (Kalita, 2011).	Essential oils, phenolic compounds, flavonoids, carbohydrates, proteins, alkaloids, glycosides, iroids glycosides, phenyl ethanoids, oligosaccharids, quinine, saponins, steroids, triterpens, sesquiterpens, tannins (Kensa, 2011; Kalita, 2011).

WP: whole plant; L: leaves; B: barks; R: roots; S: stems; SL: stems with leaves; T: tuber; F: fruits

MP: mode of preparation, are written in parenthesis for each part used; D: decoction; M: maceration; S: squeezed; I: infusion

Freq.: frequency of citation by informants

The minimum inhibition concentrations (MIC) and minimum bactericidal concentration (MBC) values are presented in Table 2. The lowest MIC values 128 µg/ml was exhibited by the extract of *Vitex doniana* against *Salmonella paratyphi* A; followed by the extract of *Rauvolfia vomitoria*, *Senna alata* and *Terminalia glaucescens* with MIC values of 512 µg/ml each on *S. paratyphi* A, while *S. paratyphi* B was most sensitive on *Senna alata* and *Terminalia glaucescens* extract with a MIC of 512 µg/ml.

The extract of *Bidens pilosa*, *Harungana madagascariensis*, *Gardenia ternifolia* and *Lantana camara* were most effective against *Salmonella typhi* with MIC values of 512 µg/ml. *Annona muricata*, *Senna alata*, *Terminalia glaucescens*,

Paullinia pinnata and *Vitex doniana* inhibited the growth of all the bacteria tested. The results of table 2 showed detectable MBC values for some of the studied plants on the tested microbial strains.

When analyzing carefully the MIC and MBC results for each plant samples, it can be noted that MBC/MIC ratios lower than 4 were obtained for *Carica papaya*, *Pseudarthria confertiflora*, *Moringa oleifera* and *Harungana madagascariensis*, thus considered to have bactericidal activities (Noumedem *et al.*, 2013). To the best of our knowledge, the activity of *Pseudarthria confertiflora* on the microorganisms studied in the present work is being reported for the first time.

Table 2: Screening of 38 plant species for potential antibacterial activity against some *Salmonella* species.

Botanic name	P Sc	<i>S. typhi</i>				SPA					
		40 mg/ml	80 mg/ml	160 mg/ml	MIC	MBC	40 mg/ml	80 mg/ml	160 mg/ml	MIC	MBC
<i>Eremomastax speciosa</i> (Hochst) cufod	L	7.67 ± 0.46	8 ± 0.81	10.5 ± 0.40	2048	4096 (2)	nd	nd	nd	1024	8192 (8)
<i>Pseudospondias microcarpa</i> Engl.	B	8.5 ± 0.40	9.5 ± 0.40	11.5 ± 0.40	4096	/	nd	nd	nd	/	/
<i>Annona muricata</i> Linn.	L	9.5 ± 0.40	11 ± 0.81	14.5 ± 0.40	4096	4096 (1)	0±0	10 ± 00	13 ± 00	2048	/
<i>Picralima nitida</i> (Stapf) Th. et H.	R	0±0	9 ± 00	12 ± 0.81	1024	4096 (4)	0±0	8.5 ± 1.22	10 ± 1.63	/	/
<i>Rauvolfia vomitoria</i> Afz.	B	0 ± 0	10 ± 00	12 ± 0.81	8192	/	11 ± 0.81	12.5 ± 0.40	15.5 ± 0.40	2048	/
<i>Dracaena deisteliana</i> Engl	L, R	0±0	0±0	7.5 ± 0.40	/	/	nd	nd	nd	/	/
<i>Bidens pilosa</i> Linn.	Wp, L	10.5 ± 1.22	12 ± 0.81	13 ± 0.81	512	2048 (4)	10 ± 00	10.5 ± 0.40	12 ± 0.81	/	/
<i>Emilia coccinea</i> (Sims) G. Don.	Wp	0±0	0±0	8.5 ± 0.41	nd	nd	0±0	0±0	8.5 ± 0.41	/	/
<i>Laggera alata</i> (G. Don) Sch. BipEx. Oliv.	L	7.5 ± 0.40	10 ± 00	13.5 ± 0.40	4096	/	nd	nd	nd	/	/
<i>Vernonia colorata</i> (Willd.) Drake	L	0 ± 00	9 ± 0.81	10.5 ± 0.40	/	/	nd	nd	nd	/	/
<i>Spathodea campanulata</i> P.Beauv.	L	9 ± 00	10 ± 0.81	12.5 ± 0.40	1024	8192 (8)	nd	nd	nd	/	/
<i>Stereospermum acuminatissimum</i> K. Schum.	B	11 ± 00	13 ± 0.81	15 ± 0.81	2048	/	nd	nd	nd	/	/
<i>Cordia platyhyrsa</i> Bak.	F	10 ± 00	11 ± 0.81	12 ± 0.81	2048	8192 (4)	nd	nd	nd	/	/
<i>Canarium Schweinfurthii</i> Engl.	B	0 ± 00	9 ± 00	10.5 ± 0.40	4096	8192 (2)	nd	nd	nd	/	/
<i>Dacryodes edulis</i> (G.Don) H.J.	L	14.5 ± 0.40	15.5 ± 0.40	19.5 ± 0.40	2048	4096 (2)	nd	nd	nd	/	/
<i>Carica papaya</i> Linn.	L, B, F	8.5 ± 0.40	10 ± 00	12 ± 0.00	1024	2048 (2)	9 ± 0.81	11 ± 0.81	12 ± 0.81	/	/
<i>Terminalia glaucescens</i> Planch.	B	9.5 ± 0.40	16.5 ± 0.40	20.5 ± 0.40	8192	/	0 ± 00	10.5 ± 0.40	13 ± 0.81	512	2048 (4)
<i>Cupressus lusitanica</i> Mill.	Wp, L, S	0 ± 00	8.5 ± 0.40	10.5 ± 0.40	2048	/	nd	nd	nd	/	/
<i>Pteridium aquilinum</i> (Linn.) kuhl.	L	0 ± 00	9 ± 0.81	11.5 ± 0.40	4096	/	0 ± 00	8.5 ± 0.40	11 ± 0.81	2048	/
<i>Dioscorea dumetorum</i>	L	0 ± 00	9.5 ± 0.40	11 ± 0.81	1024	8192 (8)	nd	nd	nd	/	/
<i>Entada abyssinica</i> A. Rich.	B	0 ± 00	8.5 ± 0.40	10.5 ± 0.40	2048	/	nd	nd	nd	/	/
<i>Pseudarthria confertiflora</i> (A. Rich.) Baker	L	11.5 ± 0.40	15 ± 0.81	23.5 ± 1.22	2048	8192 (4)	nd	nd	nd	/	/
<i>Senna alata</i> (Linn.) Roxb	L	10.5 ± 0.40	17.5 ± 0.40	20.5 ± 0.40	1024	/	12.5 ± 0.40	17.5 ± 0.40	24 ± 0.81	512	4096 (8)
<i>Harungana madagascariensis</i> Lam-expoir	B	0 ± 00	8.5 ± 0.40	11.5 ± 0.40	512	2048 (4)	nd	nd	nd	/	/
<i>Ocimum gratissimum</i> Linn	L	10.5 ± 0.40	11.5 ± 0.40	12 ± 0.40	2048	4096 (2)	0 ± 00	10.5 ± 0.40	11 ± 00	/	/
<i>Vitex doniana</i> Sweet	B	0 ± 00	0 ± 00	10 ± 00	1024	4096 (4)	9.5 ± 0.40	11 ± 0.81	11.5 ± 1.22	1024	4096 (4)
<i>Gossypium barbadense</i> Linn.	L	0 ± 00	10 ± 0.81	10.5 ± 0.40	8192	/	nd	nd	nd	/	/
<i>Theobroma cacao</i> L.	L, B	9.5 ± 0.40	10.5 ± 0.40	13.5 ± 0.40	8192	/	nd	nd	nd	/	/
<i>Thespesia populnea</i> (Linn.) soland ex corr	L	8 ± 00	10 ± 0.81	11 ± 0.81	2048	/	nd	nd	nd	/	/
<i>Ficus sur</i> Forssk	B	0 ± 00	10.5 ± 0.40	11 ± 0.00	8192	/	nd	nd	nd	/	/
<i>Moringa oleifera</i> Lam	L	9.5 ± 0.40	11 ± 00	12.5 ± 0.40	1024	2048 (2)	9.5 ± 0.40	10 ± 00	11.5 ± 0.40	/	/
<i>Musa paradisiaca</i> Linn.	L	9.5 ± 0.40	10.5 ± 1.22	12 ± 0.40	2048	4096 (2)	0 ± 00	0 ± 00	10.5 ± 0.40	/	/
<i>Psidium guajava</i> Linn.	L, B	9 ± 0.81	10 ± 00	11.5 ± 0.40	2048	4096 (2)	nd	nd	nd	/	/
<i>Imperata cylindrica</i> (L.)Raeusch.	Wp	0 ± 00	0 ± 00	10.5 ± 0.40	2048	/	nd	nd	nd	/	/
<i>Paullinia pinnata</i> Linn.	L	0 ± 00	10.5 ± 0.40x	11 ± 0.00	2048	8192 (4)	0 ± 00	9.5 ± 0.40	12 ± 0.40	2048	2048 (1)
<i>Gardenia temifolia</i> Schum et thonn.	B	0 ± 00	9.5 ± 0.40	11 ± 0.81	512	2048 (4)	nd	nd	nd	/	/
<i>Mitracarpus villosus</i> (Swartz)DC.	Wp	0 ± 00	11 ± 00	13.5 ± 0.40	4096	/	nd	nd	nd	/	/
<i>Lantana camara</i> Linn	L	10.5 ± 0.40	11.5 ± 0.40	13 ± 0.81	512	2048 (4)	8.5 ± 0.40	9 ± 0.81	11 ± 0.81	/	/
<i>Eremomastax speciosa</i> (Hochst) cufod	nd	nd	nd	/	/	-	++	++	+	+	-
<i>Pseudospondias microcarpa</i> Engl.	nd	nd	nd	/	/	-	+++	+++	+	+	+
<i>Annona muricata</i> Linn.	0±0	9±0.81	11.5 ± 0.40	4096	8192 (2)	+	+	+	+	+	+++
<i>Picralima nitida</i> (Stapf) Th. et H.	nd	nd	nd	/	/	+	+	+	+	+	+++
<i>Rauvolfia vomitoria</i> Afz.	11 ± 0.81	12.5 ± 0.40	15.5 ± 0.40	512	4096 (8)	+	+	++	+	+	-
<i>Dracaena deisteliana</i> Engl	0±0	8.5 ± 1.22	12 ± 00	/	/	+	-	-	+	+	++
<i>Bidens pilosa</i> Linn.	9 ± 00	10 ± 00	12.5 ± 0.40	/	/	+	+	+	+	+	-
<i>Emilia coccinea</i> (Sims) G. Don.	/	/	/	/	/	+	+++	+	+	+	-
<i>Laggera alata</i> (G. Don) Sch. BipEx. Oliv.	nd	nd	nd	/	/	+	+	++	+	+	+
<i>Vernonia colorata</i> (Willd.) Drake	nd	nd	nd	/	/	+	++	+++	+	+	+
<i>Spathodea campanulata</i> P.Beauv.	nd	nd	nd	/	/	+	+	+++	+	+	+
<i>Stereospermum acuminatissimum</i> K. Schum.	nd	nd	nd	/	/	+	+	+	+	+	+
<i>Cordia platyhyrsa</i> Bak.	0 ± 00	8.5 ± 0.40	9.5 ± 0.40	/	/	+	+++	+++	+	+	+++
<i>Canarium Schweinfurthii</i> Engl.	nd	nd	nd	/	/	+	+	+++	+	+	-
<i>Dacryodes edulis</i> (G.Don) H.J.	nd	nd	nd	/	/	+	+	+++	+	+	-
<i>Carica papaya</i> Linn.	0 ± 00	9.5 ± 0.40	11 ± 0.81	/	/	+	+	+	+	+	+
<i>Terminalia glaucescens</i> Planch.	nd	nd	nd	512	4096 (8)	+	+++	+++	+	+	+++
<i>Cupressus lusitanica</i> Mill.	nd	nd	nd	/	/	-	+	+	++	+	+++
<i>Pteridium aquilinum</i> (Linn.) kuhl.	0 ± 00	0 ± 00	9 ± 0.81	/	/	-	-	++	+	+	++
<i>Dioscorea dumetorum</i>	nd	nd	nd	/	/	-	-	++	-	-	++
<i>Entada abyssinica</i> A. Rich.	nd	nd	nd	/	/	-	++	++	+	+	++
<i>Pseudarthria confertiflora</i> (A. Rich.) Baker	11 ± 0.81	13.5 ± 1.22	17.5 ± 0.40	2048	2048 (1)	+	+++	+++	+	+	+++
<i>Senna alata</i> (Linn.) Roxb	10.5 ± 0.40	13.5 ± 0.40	22 ± 2.03	512	4096 (8)	+	+	++	++	+	++
<i>Harungana madagascariensis</i> Lam-expoir	nd	nd	nd	/	/	+	+++	+++	++	++	-
<i>Ocimum gratissimum</i> Linn	0 ± 00	8.5 ± 0.16	10 ± 00	/	/	+	+	+	+	+	+
<i>Vitex doniana</i> Sweet	0 ± 00	10 ± 00	11 ± 0.81	128	2048	-	++	++	+	+	+
<i>Gossypium barbadense</i> Linn.	nd	nd	nd	/	/	-	+++	+++	-	-	+
<i>Theobroma cacao</i> L.	nd	nd	nd	/	/	-	+++	+++	+	+	+
<i>Thespesia populnea</i> (Linn.) soland ex corr	nd	nd	nd	/	/	-	+	+	++	+	-
<i>Ficus sur</i> Forssk	nd	nd	nd	/	/	-	++	++	+	+	+++
<i>Moringa oleifera</i> Lam	nd	nd	nd	/	/	-	++	+++	+	+	-
<i>Musa paradisiaca</i> Linn.	0 ± 00	9 ± 0.81	12 ± 00	/	/	+	+	+	+	+	+
<i>Psidium guajava</i> Linn.	nd	nd	nd	/	/	+	+	+	+	+	+
<i>Imperata cylindrica</i> (L.)Raeusch.	nd	nd	nd	/	/	+	+	+	+	+	-
<i>Paullinia pinnata</i> Linn.	8.5 ± 1.22	10.5 ± 1.2	13 ± 0.81	1024	8192 (8)	+	-	+	+	+	+++
<i>Gardenia temifolia</i> Schum et thonn.	nd	nd	nd	/	/	-	+++	+++	+	+	+
<i>Mitracarpus villosus</i> (Swartz)DC.	nd	nd	nd	/	/	-	+++	+++	+	+	+++
<i>Lantana camara</i> Linn	9.5 ± 0.40	11 ± 0.81	12.5 ± 1.22	8192	/	-	+	+	+	+	-

	I.Z.D	MIC	MBC	MBC/MIC	
Ciprofloxacin (10 µg/ml)	30 ± 0.81	64	128	2	<i>S. typhi</i> ATCC 6539
	28.5 ± 1.22	32	64	2	SPA
	30 ± 1.63	32	64	2	SPB

P Sc: part screened., I.Z.D.= Mean diameter of growth inhibition zones in mm average of three replicates, SD= Standard deviation. *S. typhi* : *Salmonella typhi* ; SPA: *Salmonella paratyphi A* ; SPB : *Salmonella paratyphi B.*, Al: Alkaloids; Fl: Flavonoids; Pl: Polyphenols; T: Triterpens ; Se : Sterols ; Sa : Saponins., /: indicates no zone of inhibition or activity; the result are mean ±S.D (n=3), +++, high concentration; ++, medium concentration; +, low concentration; -, not detected

DISCUSSION

The high diversity of species used for treatment of typhoid fever in the study area, is an indication of the importance of medicinal plants. The reliance on traditional medicine may be due to the perceived potency of the plants. High consensus among users in different countries reflects the significance of medicinal plants to the people (Cotton, 1996). It was clear that the informants harvest the bark, roots, stem, fruits and leaves; but the part that is used most was found to be the leaves. Leaves as frequently used organ in traditional herbal drugs is also reported elsewhere (Muthu *et al.*, 2006; Panghal *et al.*, 2010). In addition to this, leaves are the main photosynthetic organs in plants and are considered to be the natural pharmacy for synthesis of many active constituents those are pharmacologically more active against certain diseases (Passalacqua *et al.*, 2007). In general the use of leaves as the chosen plant part is a more sustainable practice as opposed to where roots and/or the bark are used. The prevalence in the use of leaves for preparation of traditional herbal remedies has been reported in other studies (Muthu *et al.* 2006; Focho *et al.* 2009; Panghal *et al.* 2010; Amri & Kisangau 2012). This practice helps to increase the chances of species survival and enhances the sustainable management of plants, as long as only an appreciable amount of leaves is harvested (Abede & Ayehu, 1993; Tadesse *et al.* 2005).

It was not surprising that decoction was the preferred preparation method. Decoction is known to be an effective extraction method compared to maceration in cold water, since boiling facilitates the extraction of constituents and also kills microorganisms potentially associated with the harvested plants. Plant parts were generally prepared using water as the solvent, likely because water is a readily available and cheap solvent and provides good solubility of the active components. Oral administration was the only mode of dispensing of herbal medicines against typhoid fever. This is because the causal bacterium is located in the intestinal tract. This mode of administration of herbal medicine was also reported elsewhere (Kamatenesi & Oryem-Origa, 2006; Bhattarai *et al.* 2010).

Bidens pilosa is the most commonly used plant for treatment by informant. Several studies document good antimicrobial effects of its extract (Rojas *et al.*, 2006; Deba *et al.*, 2008).

Phytochemical screening conducted on the plant extracts (Table 2) revealed the presence of compounds which are known to exhibit biological as well as physiological activities (Sofowora, 1993). These are alkaloids, flavonoids, polyphenols, saponins, sterols and triterpenes. The presence of all the secondary metabolites in *Annona muricata*, *Lagdera alata*, *Spathodea campanulata*, *Cordia platythirsa*, *Carica papaya*, *Terminalia glaucescens* and *Pseudarthria confertiflora*, is an indication that these plants are of pharmacological importance (Adebayo & Ishola, 2009), and also justified their potential use as drug in the study area. Almost all the plants contained triterpenes except in *Dioscorea dumetorum* and *Gossypium barbadense*. Triterpenes

have been found to be useful in the prevention and therapy of several diseases, including cancer. Triterpenoids are also known to possess antimicrobial, antifungal, antiparasitic, antiviral, anti-allergenic, antispasmodic, antihyperglycemic, antiinflammatory and immunomodulatory properties (Wagner & Elmadfa, 2003). The presence of this compound probably justified the use of the selected plants for the treatment of typhoid fever. However, the absence of this constituent in the leaves of *Gossypium barbadense* did not corroborates the findings of Charu *et al.*, (2012).

Furthermore, most of these plants contained phenolic compounds (Polyphenols and flavonoids) which are one of the largest and most ubiquitous groups of plant metabolites (Singh *et al.*, 2007). They possess biological properties such as antiapoptosis, antiaging, anticarcinogenesis, antiinflammation, antiatherosclerosis, cardiovascular protection, improvement of endothelial function, as well as inhibition of angiogenesis and cell proliferation activities (Han *et al.*, 2007). Flavonoids are hydroxylated phenolic substances known to be synthesized by plants in response to microbial infection and they have been found to be antimicrobial substances against wide range of microorganisms *in vitro*. Their activity is probably due to their ability to complex with extracellular and soluble proteins and to complex with bacterial cell wall (Marjorie, 1996). In addition, Polyphenols have been reported to exhibit antibacterial activities with distinguished characteristics in their reactivity with proteins related polyamides polymers (Haslam, 1996).

Sterols present in most of our plant sample, have been reported to have antibacterial properties (Raquel, 2007) and they are very important compounds especially due to their relationship with compounds such as sex hormones (Okwu, 2001).

The absence of alkaloids and saponins in *Moringa oleifera* in the present work is in contrast with the opinion of Gills (1992) and Kasolo *et al.*, (2010) who noted that saponins and alkaloids are two of the active constituents. For instance, alkaloids are nitrogen-containing naturally occurring compound, commonly found to have antimicrobial properties due to their ability to intercalate with DNA of the microorganisms. Also, the presence of saponin in *Annona muricata* is in agreement with previous findings of Yusha'a *et al.* (2011).

The activity of plant extracts against bacteria have been studied for years, but in more intensified way during the last three decades. During this period, numerous antimicrobial screening evaluations have been published based on the traditional Chinese, African and Asian uses of plant-based drugs (Suffredim *et al.*, 2004). In the present study, the results of antibacterial property of the 38 plant extracts (table 2) against tested organisms varied depending on bacteria tested and concentration as previously reported by (Ravikumar *et al.*, 2007; Rajesk *et al.*, 2010). However, 38, 15 and 14 plants shows inhibition diameter against *Salmonella typhi*, *S. paratyphi B* and *S. paratyphi A* respectively. Nevertheless, negative results obtain with some of the plants extracts tested do not indicate the absence of bioactive constituents, nor that the plant is inactive. Active compound(s) may be present in insufficient quantities in the extracts to show

activity with the dose levels employed (Taylor *et al.*, 2001). Lack of the activity can thus only be proven by using large dose (Farnsworth, 1993), if the active principle is present in high enough quantities, there could be other constituents exerting antagonistic effects or negating the positive effects of the bioactive agents (Jager *et al.*, 1999).

It was noticed that *Senna alata*, showing the highest antibacterial activity on both pathogens with zone of inhibition of 24, 22.5, 20.5 mm on the *Salmonella paratyphi A*, *S. paratyphi B* and *S. typhi* respectively at 160 mg/ml. This could probably have been due to the fact that the rate of the active ingredients or constituents in the plant materials is higher compared to the other plants used in this research (Kunle & Egharevba, 2009). Preliminary phytochemical analysis of *S. alata* leaves showed that, they possess polyphenols, triterpenes and saponins. Phytoconstituents such as polyphenols, triterpenes and saponins have been reported to inhibit bacterial growth and to be protective to plants against bacterial and fungal infections (Mather & Gonzalel, 1982; Okwute, 1992). The presence of saponins in this plant must have exhibited direct antibacterial activity and suppression of bacterial virulence resulting to the antimicrobial activity seen in this study (Gills, 1992). This also corroborate with the report of Owoyale *et al.*, (2005) who documented on the plant *Senna alata* to be effective in treatment of fungal and bacterial diseases.

The fact that *Salmonella typhi* was more susceptible to the extract of *Pseudarthria confertiflora*, *Terminalia glaucescens*, *Senna alata*, *Dacryodes edulis* and *Stereospermum acuminatissimum* indicated the potency of these plants against typhoid fever. *Senna alata* and *Rauvolfia vomitoria* were found to be the most sensitive against *Salmonella paratyphi A*. However, *Salmonella paratyphi B* was sensitive to the extract of *Senna alata*, *Pseudarthria confertiflora* and *Rauvolfia vomitoria*. The antibacterial activity of *Terminalia glaucescens*, *Senna alata*, *Dacryodes edulis* and *Rauvolfia vomitoria* have been reported in the literature (Owoyale *et al.*, 2005; Obame *et al.*, 2008; Adebayo and Ishola, 2009; Ayepola, 2009; Ogundiya *et al.*, 2009; Ajibesin *et al.*, 2011; Bolou *et al.*, 2011; Omogbai & Eneh, 2011).

In local area, these plants are being used in the treatment of typhoid fever, diarrhea, dysentery etc. So, the uses of these plants as antibacterial agent are justified; since they showed good inhibition zones. Demonstration of low MIC (128 µg/ml) especially by *Vitex doniana* (table 2) is an indication that the phytoconstituents of the plant have therapeutic potential. Similar report revealed the antimicrobial potency of *Vitex doniana* with MIC range from 0.4-128 µg/ml (Iroha *et al.*, 2010). It is important to highlight that *Vitex doniana* has been thoroughly studied phytochemically and its antimicrobial activity has been evaluated (Kubmarawa *et al.*, 2007; Iroha *et al.*, 2010). In this study, its antibacterial activity was demonstrated. It is no surprising the activity shown by this plant, since other species of the same genus (*V. negundo*) have shown to have antibacterial, anti-inflammatory and anti-fungal activity (Rusia & Srivastava, 1998; Dharmasiri *et al.*, 2003; Panda *et al.*, 2009).

Based on literature on these 38 plants, there is no biological and chemical on *Pseudarthria confertiflora*. Several species of the genus *Pseudarthria* (*P. viscida* and *P. hookeri*) has been studied and it has been demonstrated that they presented immunomodulatory and anti-fungal activity (Clarkson *et al.*, 2004; Deepa *et al.*, 2004; Mathew & Sasikumar, 2007; Vijayabaskaran *et al.*, 2010), but the particular species *Pseudarthria confertiflora* has never been studied before and in this work, its antibacterial activity was demonstrated. The activity of this plant against *salmonella typhi*, *S. paratyphi A* and B has added one more plant that can offer alternative medicare to diseases caused by these microorganisms.

According to Noumedem *et al.* (2013), a sample is bactericidal when the ratio MBC/MIC ≤ 4 and bacteriostatic when this ratio is >4 . It therefore appeared that bactericidal effects were obtained with the extracts from *Carica papaya*, *Pseudarthria confertiflora*, *Moringa oleifera* and *Harungana madagascariensis*. The result of antibacterial of *Vitex doniana*, *Rauvolfia vomitoria*, *Senna alata*, *Terminalia glaucescens* *Bidens pilosa*, *Harungana madagascariensis* *Gardenia ternifolia* and *Lantana camara* is highly encouraging and gives new lead plants for isolation and characterization of the active compounds. It is probably these active chemical constituents present in these plants which are responsible for the treatment of various bacterial infections caused by *Salmonella sp.*

CONCLUSION

The result of the present study offers a scientific basic for the use of the majority of tested plants in the treatment of typhoid fever in Bamboutos division. These plants species can be regarded as promising resources for anti-typhoid drugs. It seems that further investigations are necessary in order to draw solid conclusions.

REFERENCES

- Abdulrahma F. 1992. Studies on natural products. The Moraceae in African traditional medicine and management of psychiatry in Borno State. M.Sc. Thesis, University of Maiduguri, Maiduguri, Pp.89-95.
- Abebe D, Ayehu A. 1993. *Medicinal Plants and Enigmatic Health Practices in Northern Ethiopia*. Berhanina Selam Printing Enterprise, Addis Ababa, Ethiopia.
- Abubakar BA, Aliyu MM, Mikhail SA, Hamisu I, Adebayo O. Phytochemical screening and antibacterial activities of *Vernonia ambigua*, *Vernonia blumeoides* and *Vernonia ocephala* (Asteraceae). *Natural Drugs*, 2009.
- Addo-Fordjour P, Kofi Anning A, Durosimi Belford E J, Akonnor D. Diversity and conservation of medicinal plants in the Bomaa community of the Brong Ahafo Region, Ghana. *Journal of Medicinal Plants Research*, 2008; 2(9): 226-233.
- Adebayo EA, Ishola OR. Phytochemical and antimicrobial screening of crude extracts from the root, stem bark, and leaves of *Terminalia glaucescens*. *African Journal of Pharmacy and Pharmacology*, 2009; 3(5):217-221.
- Adeneye AA, Olagunju JA, Elias SO, Olatunbosun DO, Mustafa AO, Adeshile OI, Ashaolu AO, Laoye TA, Bamigboye AO, Adeoye AO. Protective activities of the aqueous root extract of *Harungana madagascariensis* in acute and repeated acetaminophen

hepatotoxic rats. *International Journal of Applied Research in Natural Products*, 2008; 1(3): 29-42.

Adinortey MB, Sarfo JK, Aduko GE, Dzotsi E, Kusi S, Ahmed MA, Abdul-Gafaru O. Acute and sub-acute oral toxicity assessment of hydro-alcoholic root extract of *Paullinia pinnata* on haematological and biochemical parameters. *Biology and Medicine*, 2012; 4(3): 121-125.

Adjanohoun EJ, Aboubakar N, Dramane K, Ebot ME, Ekpere JA, Enow-Orock EG, Focho D, Gbile ZO, Kamanyi A, Kamsukom J, Keita A, Mbeukum I, Mbi CN, Mbiele AL, Mbome LL, Mobiru NK, Nancy WL, Nkongmeneck B, Sofowora A, Tamze V, Wirmum CK. 1996. Traditional medicine and pharmacopoeia. Contribution to ethnobotanical and floristic studies in Cameroon. *Organisation of African unity. Scientific, Technical and research commission* (OAU/STRC) Lagos, Nigeria; 636p.

Agyare C, Asase A, Lechtenberg M, Nihues M, Deters A, Hensel A. An ethnopharmacological use of medicinal plants used for wound healing in Bosomtwi-Atwima-Kwanwoma area, Ghana. *Journal of Ethnopharmacology*, 2009; 125: 393-403.

Ajaiyeoba EO, Abiodun OO, Falade MO, Ogbale NO, Ashidi JS, Happi CT, Akinboye DO. *In vitro* cytotoxicity studies of 20 plants used in nigerian antimalarial ethnomedicine. *Phytomedicine*, 2006; 13: 295-298.

Ajibesin KK, Ekpo BA, Bala DN, Essien EE, Adesanya SA. Ethnobotanical survey of Akwa Ibom State of Nigeria. *Journal of Ethnopharmacology*, 2008; 115: 387-408.

Ajibesin KK, Essien EE, Adesanya SA. Antibacterial constituents of the leaves of *Dacryodes edulis*. *African Journal of Pharmacy and Pharmacology*, 2011; 5(15):1782-1786.

Akila A, Rani K. Biosynthesis of gossypol in *Thespesia populnea*. *Phytochemistry*, 1993; 335-340.

Amri E, Kisangau DP. Ethnomedicinal study of plants used in villages around Kimboza forest reserve in Morogoro, Tanzania. *Journal of Ethnobiology and Ethnomedicine*, 2012; 8:1. <http://dx.doi.org/10.1186/1746-4269-8-1>.

Amusan OOG, Adesogan EK, Makinde, JM. Antimalarial active principles of *Spathodea campanulata* stem bark. *Phytotherapy Research*, 1996; 10(8):692-693.

Amusan OOG, Msonthi JD, Makhubu LP. Molluscicidal activity of *Spathodea campanulata*, *Andrachne ovalis*, *Phytolacca dodecandra* and *Hypoxis rooperi*. *Fitoterapia*, 1995; 66:113-116.

Anbu JSJS, Anandarajagopal K, Dinesh KC, Rejitha G, Suraj R, Hj Azman H, Vignesh M, Proom P. Ethnomedical survey of plants used by the Orang Asli in Kampung Bawong, Perak, West Malaysia. *Journal of Ethnobiology and Ethnomedicine*, 2010; 6:5.

Annan K, Houghton PJ. Two novel lupine triterpenoids from *Paullinia pinnata* L. with fibroblast stimulatory activity. *Journal of Pharmacy and Pharmacology*, 2010; 62(5): 663-668.

Arbonnier M. 2002. Arbres, arbustes et lianes des zones sèches d'Afrique de l'Ouest. CIRAD Maseum national d'histoire naturelle, 573p.

Asfaw N, Storesund HJ, Aasen AJ. Constituents of the essential oil of *Laggera tomentosa* Sch. Bip. ex Oliv. et Hiern endemic to Ethiopia. *J. Essent. Oil Res.* 2003, 15, 102-106.

Atta-ur-Rahman, Choudhary MI, Thomsen WJ. 2005. Bioassay Techniques for Drug Development. Taylor & Francis e-Library, Amsterdam, the Netherlands.

Awal MA, Aina Nahar M, Shamin Hossain MA, Barri M, Rahman Haque ME. Brine shrimp toxicity of leaf and seed extracts of *Cassia alata* Linn. and their antibacterial potency. *J. Med. Sci.* 2004; 4(3): 188-193.

Ayepola OO. Evaluation of the antimicrobial activity of root and leaf extracts of *Terminalia glaucescens*. *Advances in Natural and Applied Sciences*, 2009; 3(2):188-191.

Ayuk ET, Duguma B, Franzel S, Kangué J, Mollet M, Tiki-Manga T, Zekeng, T. Uses, management and economic potential of *Dacryodes edulis* (Burseraceae) in the humid low lands of Cameroon. *Econ. Bot.* 1999; 53(3): 292-301.

Balancod DT, Vallejo LV, Patacsil M, Apostol O, Laruan AVML, Manuel J, Cortez S, Gutierrez, MR. Phytochemical screening and

antibacterial activity of selected medicinal plants of Bayabas, Sablan, Benguet Province, Cordillera Administrative Region, Luzon, Philippines. *Indian Journal of Traditional Knowledge*, 2012; 11(4): 580-585.

Banerjee A, De, B. Anthocyanins in some flowers of West Bengal. *J Med Arom Plant Sci.*, 2001; 23: 600-604.

Baskar R, Rajeswari V, Sathish Kumar T. *In vitro* antioxidant studies in leaves of *Annona* species. *Indian Journal of Experimental Biology*, 2007; 45:480-485.

Bekele-Tesemma A, Birnie A, Tengnas B. 1993. Arbres et arbustes utiles pour l'Ethiopie. Unité de conservation des sols régional (RSCU), l'Agence suédoise de développement international (SIDA).

Békro YA, Békro JAM, Boua BB, Trab FH, Ehilé EE. Etude ethnobotanique et screening phytochimique de *Caesalpinia benthamiana* (Baill.) Herend. Et Zarucchi (Caesalpinaceae). *Rev. Sci. Nat.*, 2007; 4(2): 217-225.

Bellomaria B, Kacou P. plantes et médecine populaire d'Agboville (Côte d'Ivoire). *Fitoterapia*, 1995; 66: 117.

Benoit F, Valentin A, Pelissier Y, Diafouka F, Marion C. *In vitro* antimalarial activity of vegetal extracts used in West African traditional medicine. *American Journal of Tropical Medicine and Hygiene*, 1996; 54(1): 67-71.

Benoit VF, Santillama H, Kone BD. Anti-toxoplasma activity of vegetal extracts used in West African traditional medicine. *Parite*, 2000; 7(1): 3-7.

Bharali R, Tabassum J, Azad MRH. Chemomodulatory effect of *Moringa oleifera* Lam. on hepatic carcinogen metabolizing enzymes, antioxidant parameters and skin papillomagenesis in mice. *Asian Pacific Journal of Cancer Prevention*, 2003; 4: 131-139.

Bhattarai S, Chaudhary RP, Quave CL, Taylor RSL. The use of medicinal plant species in the trans-himalayan arid zone of Mustang district, Nepal. *Journal of Ethnobiology and Ethnomedicine*, 2010; 6:14. <http://dx.doi.org/10.1186/1746-4269-6-14>.

Biswas KR, Ishika T, Rahman M, Khan T, Swarna A, Monalisa MN, Sanam S, Malek I, Rahmatullah M. Medicinal plants used for preventive medicinal purposes: a survey in Muktipara village, Chuadanga district, Bangladesh. *American Eurasian Journal of Sustainable Agriculture*, 2011; 5: 247-251.

Bolou GEK, Attioua B, N'Guessan AC, Coulibaly A, N'Guessan JD, Djaman AJ. Évaluation *in vitro* de l'activité antibactérienne des extraits de *Terminalia glaucescens* planch. sur *Salmonella typhi* et *Salmonella typhimurium*. *Bulletin de la Société Royale des Sciences de Liège*, 2011; 80: 772-790.

Bouquet TA. 1969. Féticheurs et Médecines traditionnelles du Congo-Brazzaville, Mémoires O.R.S.T.O.M. no 36,1 vol. 306 p., Fig., Pl.,

Brandão MGL, Kretli AU, Soares LSR, Nery CGC, Marinuzzi HC. Antimalarial activity of extracts and fractions from *Bidens pilosa* and other *Bidens* species (Asteraceae) correlated with the presence of acetylene and flavonoid compounds. *J. Ethnopharmacol.* 1997; 57: 131-138.

Burkill HM. 1985. The useful plants of west tropical Africa. Royal Botanic Gardens Kew, vol.1, 2nd Ed. P495.

Charu AC, Sanjeev M, Heena D. Phytochemical Screening and Evaluation of Biological Activities of Some Medicinal Plants of Phagwara, Punjab. *Asian Journal of Chemistry*, 2012; 24(12): 5903-5905.

Chiang YM, Chang CLT, Chang SL, Yang WC, Shyur LF. Cytopylone, a novel polyacetylenic glucoside from *Bidens pilosa*, functions as a T helper cell modulator. *J. Ethnopharmacol.* 2007; 110: 532-538.

Chifundera K. Livestock diseases and the traditional medicine in the Bushi area, Kivu province, Democratic Republic of Congo. *African Study Monographs*, 1998; 19(1): 13-33.

Clarkson C, Maharaj VJ, Crouch NR, Grace OM, pillay P, Matsabisa MG, Bhagwandin N, Smith PJ, Folb PI. *In vitro* antiplasmodial activity of medicinal native to or naturalised in South Africa. *Journal of Ethnopharmacology*, 2004; 92:177-191.

Cos P, Hermans N, De Bruyne T, Apers S, Sindambiwe JB, Witvrouw M, De Clercq E, Vanden Berghe D, Pieters L, Vlietinck AJ. Antiviral activity of rwandan medicinal plants against human Immuno-

- deficiency virus type 1 (HIV-1). *Phytomedicine*, 2002; 9: 62-68.
- Cotton CM. 1996. *Ethnobotany: Principles and Applications*. Chichester: John Wiley and Sons Ltd.
- Daziel JM. 1937. *The useful plants of West Africa: Appendix to the flora of West Tropical Africa*. The Crown Agents for the Colonies: London; 612p.
- Deba F, Xuan TD, Yasuda M, Tawata S. Chemical composition and antioxidant, antibacterial and antifungal activities of the essential oils from *Bidens Pilosa* Linn. *Var. Radiata*. *Food Control*, 2008; 19: 346-352.
- Dharmasiri MG, Jayakody JRAC, Galhena G, Liyanage SSP, Ratnasooriya WD. Anti-inflammatory and analgesic activities of mature fresh leaves of *Vitex negundo*. *J. Ethnopharmacol*, 2003; 87: 199-206
- Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology*, 2005; 4(7): 685-688.
- Ekpendu TE. 1995. A study of the extractives of three medicinal plants. Ph.D. Thesis, University of Ibadan, 47p.
- Ekundayo O, Oguntimein B, Laakso I, Hiltunen R. Composition of the essential oil of *Laggera alata*. *Planta Med*, 1989; 55: 573-574.
- Elujoba AA. Studies on the antiarrhoea activity of *Ocimum gratissimum*. *University of Ife press*, 2000.
- Elusiyan CA, Ani NC, Adewunmi CO, Olugbade TA. Distribution of iridiod glucosides and anti-oxidant compounds in *Spathodea campanulata*. *Afr J Tradit Complement Altern Med*, 2011; 8(1): 27-33.
- Fabry W, Paul O, Rainer A. Antibacterial activity of East African Medicinal. *Journal of Ethnopharmacology*, 1998; 60: 79-84.
- Fahey JW. *Moringa Oleifera*: A Review of the Medical evidence for its Nutritional, Therapeutic and prophylactic properties. *Trees life J*, 2005; 15(1): 1-15.
- Faleyimi OI, Okinyemi O, Idris YM. Survey of forest plants used in traditional treatment of typhoid fever in Chikun local government area of Kaduna state, Nigeria. *International Journal of Biomedical and Health Sciences*, 2010; 6(1): 75-81.
- Farnsworth NR. 1993. Biological approaches to the screening and evaluation of natural products. In: Rasoanaivo P, Ratsimamanga-Urverg S (Eds) *Biological evaluation of plants with reference to the Malagasy Flora, Madagascar*, pp. 35-43.
- Focho DA, Nkeng EAP, Lucha CF, Ndam WT, Afegenui A. Ethnobotanical survey of plants used to treat diseases of the reproductive system and preliminary phytochemical screening of some species of Malvaceae in Ndop Central Subdivision, Cameroon. *Journal of Medicinal Plants Research*, 2009; 3(4): 301-314.
- Freiburghaus F, Steck A, Pfander H, Burn R. 1998. Bioassay-guided isolation of diastereoisomer of kolavenol from *Entada abyssinica* active on *trypanosome brucei rhodesiense*. *Journal of Ethnopharmacology*, 1998; 61 (3): 179-183.
- Gasquet, Bamba. 1985. Action amoebicide et anthelminthique du vernolide et de l'hydroxyvermolide isolés des feuilles de *Vernonia colorata* (Willd) Drake. *European Journal of Medicinal Chemistry*, 1985 ; 20: 111-115.
- Ghani A. 2003. Medicinal plants of Bangladesh: Chemical constituents and uses. 2nd Ed. *The Asiatic Society of Bangladesh*, Dhaka Bangladesh, 315p.
- Gills LS. 1992. *Ethnomedical uses of plants in Nigeria*. University of Benin press, Nigeria. P. 276.
- Gurib-Fakim A. 2008. *Toutes les plantes qui soignent. Plantes d'hier, médicaments d'aujourd'hui*. Editions Michel Lafon, 734p.
- Han X, Shen T, Lou H. Dietary polyphenols and their biological significance. *Int. J. Mol. Sci*, 2007; 950-988.
- Hashim H, Kamali EL, Yagoub El-Amir M. Antibacterial activity and phytochemical screening of ethanolic extracts obtained from selected Sudanese medicinal plants. *Curr. Res. J. Biol Sci*, 2010; 2(2): 143-146.
- Haslam E. Natural polyphenols (vegetable tannins) as drugs: possible mode of action. *J. Natl. Prod*, 1996; 59:205-215.
- Hassan A, Khan MA, Ahmad M. Authenticity of folk medicinal plants of Pakistan. *Taxonomic Chemical Methods*, 2007; 1: 1-5.
- Hegnauer R. 1973. *Chemotaxonomie der pflanzen*, Birkhäuser Verlag, Basel, Stuttgart, 6, 761p.
- Imaga NA, Gbenle G, Okochi VI, Adenekan S, Duro-Emmanuel T, Oyeniyi B, Dokai PN, Oyenuga M, Otumara A, Ekeh FC. 2010. Phytochemical and antioxidant nutrient constituents of *Carica papaya* and *Parquetina nigrescens* extracts. *Scientific Research and Essays*, 2010; 5(16): 2201-2205.
- Iroha IR, Ilang DC, Ayogu TE, Oji AE, Ugbo EC. Screening for anti-typhoid activity of some medicinal plants used in traditional medicine in Ebonyi State Nigeria. *African Journal of Pharmacy and Pharmacology*, 2010; 4(12):860-864.
- Jager AK, Hutchings A, Van Staden J. Screening of plants used by southern African traditional healers in the treatment of dysmenorrhoea for prostaglandin-synthesis inhibitors and uterine relaxing activity. *J. Ethnopharmacol*, 1999; 64:9-14.
- Jager AK, Hutchings A, Van Staden J. Screening of Zulu medicinal plants for prostaglandin-synthesis inhibitors. *J. Ethnopharmacol*, 1996; 52: 95-100.
- Jegade IA, Kunle OF, Ibrahim JA, Ugbabe G, Okogun JI. Pharmacognostic investigation of leaves of *Mitracarpus vilosus* (S.W.) DC. *African Journal of Biotechnology*, 2005; 4(9): 957-959.
- Kamatenesi MM, Acipa A, Oryem-origa H. Medicinal plants of Otwal and Ngai Sub counties in Oyam District, Northern Uganda. *Journal of Ethnobiology and Ethnomedicine*, 2011; 7:7
- Kamatenesi-Mugisha M, Oryem-Origa H. Medicinal plants used to induce labour during childbirth in western Uganda. *Journal of Ethnopharmacology*, 2006; 109:1-9. <http://dx.doi.org/10.1016/j.jep.2006.06.011>.
- Kasolo J, Bimenya SG, Ojok L, Ochieng J, Ogwal-Okeng WJ. Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*, 2010; 4(9):753-757.
- Kelmanson JE, Jäger AK Van Staden J. Zulu medicinal plant with antibacterial activity. *Journal of Ethnopharmacology*, 2000; 69: 241-246.
- Kerharo J, Adam JG. 1974. *La pharmacopée sénégalaise traditionnelle. Plantes médicinales et toxiques, Vigot frères, Paris*, 1007p.
- Khan MR, Kihara M, Omoloso AD. Anti-microbial activity of *Bidens pilosa*, *Bischofia javanica*, *Elmerillia Papuana* and *Sigesbekia orientalis*. *Fitoterapia*, 2001; 72: 662-665.
- Kolapo AL, Ogundiya MO, Okunade MB, Adejumobi JA. Assessment of phytochemical composition and Antimicrobial activity of *Terminalia glaucescens* against some oral pathogens. *EJEAFChe*, 2009; 8 (7): 466-471.
- Kouam SF, Ngadjui BT, Krohn K, Wafo P, Ajaz A, Choudhary MI. Prenylated anthronoid antioxidants from the stem bark of *Harungana madagascariensis*. *Brazilian Journal of Medical and Biological Research*, 2006:38, 1087-1094.
- Kouam SF, Yapna DB, Krohn K, Ngadjui BT, Ngoupayo J, Choudhary MI, Schulz B. Antimicrobial prenylated anthracene derivatives from the leaves of *Harungana madagascariensis*. *Journal of Natural Products*, 2007; 70, 600-603.
- Koudou J, Abena AA, Ngaissona P, Bessière JM. Chemical composition and pharmacological activity of essential oil of *Canarium schweinfurtii*. *Fitoterapia*, 2005; 76: 700-703.
- Kubmarawa D, Ajoku GA, Enwerem NM, Okorie DA. Preliminary phytochemical and antimicrobial screening of 50 medicinal plants from Nigeria. *African Journal of Biotechnology*, 2007; 6(14):1690-1696.
- Kuete V. Potential of cameroonian plants and derived products against microbial infections: *Planta Medica*, 2010; 76:1-13.
- Kunle OF, Egharevba HO. Preliminary studies on *Vernonia ambigua*: Phytochemistry and Antimicrobial Screening of the Whole Plant. *Ethnobotanical Leaflets*. 2009;13: 1216-21.
- Leume S. Multidrug resistant salmonella enteric serotype typhimurium associated with pest and Rodent. *Nein England Journal of medicine*, 1999; 356 (1): 21-28.

- Li, XC, Huo CH, Shi QW, Kiyota H. Chemical constituents of the plants from the genus *Laggera*. *Chem. Biodevers.* 2007, 4 :105–111.
- Marjorie C. Plant products as antimicrobial agents. *Clinical Microbiol. Rev.* 1996; 12:564-582.
- Martin, G.J. 1995. *Ethnobotany. A methods manual*. Chapman & Hall, London.
- Mather SB, Gonzalel L. Identification of terpenoids from leaves of *Piptocarpha peritoria* and their biological activities. *J. Nat. Prod.* 1982; 45:495-496.
- Mayer G. Antibiotics, protein synthesis, nuclei acid synthesis and metabolism. Bacteriology-chapter six. *Microbiology and Immunology on-line*, 2007; 1-11.
- Mboso OE, Eyong EU, Odey MO, Osakwe E. Comparative phytochemical screening of *Eremomastax speciosa* and *Eremomastax polysperma*. *J. Nat. Prod. Plant Resour.* 2013; 3 (2):37-41.
- Mesfin T, Debela H, Yehenew G. Survey of medicinal plants used to treat human diseases in Seka Chekorsa, Jimma zone, Ethiopia. *Ethiop. J. Health Sci.* 2005; 15(2): 89-107.
- Miller SI, Hohmann EL, Pegues DA. 1994. *Salmonella* (including *Salmonella typhi*). In principles and practice of infectious diseases Vol. 2, 4th ed. Edited by: Mandell, G.L., Bennet, J.R., Dolin, R. New York: Livingstone, pp. 2013-2033.
- MINEF (Ministry of Environment and Forestry). 1999. *Annual Report of Activities Carried Out by the Divisional Section for Forestry, Bamboutos Division*. Ministry of Environment and forestry Cameroon.
- Morton JF. 1987. Banana. In: Fruits of warm climates. Florida Flair Books, Miami.
- Moulari B, PellequerY, Lboutounne H, Girard C, Chaumont JP, Millet J, Muyard F. Isolation and in vitro antibacterial activity of astilbin, the bioactive flavanone from the leaves of *Harungana madagascariensis* Lam. ex Poir. (Hypericaceae) *Journal of Ethnopharmacology*, 2006 ; 106, 272–278.
- Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *J Ethnobiol Ethnomed.* 2006; 2: 43. doi: 10.1186/1746-4269-2-43.
- N'Guessan K, Kadja B, Zirih GN, Traoré D, Aké-Assi L. Screening phytochimique de quelques plantes médicinales ivoiriennes utilisées en pays Kroubou (Agboville, Côte d'Ivoire). *Science et Nature*, 2009 ; 6(1) : 1-15.
- Nagla'a MN. Phytochemical and antioxidant activity of *Spathodea campanulata* P. Beauvois. growing in Egypt. *Natural Product Sciences*, 2007; 13(1):11-16.
- Nanfack NAC, Fonteh FA, Payne VK, Katte B, Fogoh JM. 2014. Eau non conventionnelles: Un risque ou une solution aux problèmes d'eau pour les classes pauvres. *Larhyss Journal*, 2014; 17:47-64.
- Ndjakou Lenta B, Ngouela S, Fekam Boyom F, Tantangmo F, Feuya Tchouya GR, Tsamo E, Gut J, Rosenthal PJ, Donald Connolly J. Anti-plasmodial activity of some constituents of the root bark of *Harungana madagascariensis* LAM. (Hypericaceae). *Chemical and Pharmaceutical Bulletin*, 2007; 55:464–467.
- Ndukwe KC, Okeke IN, Lamikanra A, Adesina SK, Aboderin O. Antibacterial Activity of aqueous extract of selected chewing sticks. *Journal of Contemporary Dental Practice*, 2005; 3(6): 86-94.
- Niyonzima G, Laekeman G, Witvrouw M, Van Poel B, Pieters L, Paper D. Hypoglycemic, anticomplement and anti- HIV activities of *Spathodea campanulata* stem bark. *Phytomedicine*, 1999; 6: 45-49.
- Njomen BSN, Kamgang R, Soua P, Oyono J, Njikam N. Protective Effect of MethanolMethylene Chloride Extract of *Terminalia glaucescens* Leaves on Streptozotocin-Induced Diabetes in Mice. *Trop. J. Pharm. Res.* 2009; 8(1): 19-26.
- Noumedem J, Mihasan M, Lacmata S, Stefan M, Kuate J, Kuete V. Antibacterial activities of the methanol extracts of ten Cameroonian vegetables against Gram negative multidrug-resistant bacteria. *BMC Complement Altern Med.* 2013; 13:26.
- Obame LC, Edou P, Bassole IHN, Koudou J, Agnanié H, Eba F, Traore AS. Chemical composition, antioxidant and antimicrobial properties of the essential oil of *Dacryodes edulis* (G.Don)H.J. Lam from Gabon. *African Journal of Microbiology Research*, 2008; 2:148-152.
- Obame LC, Koudou J, Kumulungui BC, Bassole IHN, Edou P, Ouattara AS, Traore AS. Antioxidant and antimicrobial activities of *Canarium schweinfurthii* Engl. Essential oil from Centrafrican Republic. *African Journal of Biotechnology*, 2007; 6 (20): 2319-2323.
- Oben JE, Assi SE, Agbor GA, Musoro DF. Effect of *Eremomastax speciosa* on experimental diarrhea. *Afr. J. Trad. Cam.* 2006; 3: 95-100.
- Ogundiya MO, Kolapo AL, Okunade MB, Adejumbi JA. Assessment of phytochemical composition and antimicrobial activity of *Terminalia glaucescens* against some oral pathogens. *EJFAFChE*, 2009; 8(7):466-471.
- Ojewole JA. 2005. Methods find. *Exp. Clin. Pharmacol.*, 27: 547-554.
- Okafor J, Ham R. Identification, utilisation et conservation des plantes médicinales dans le Sud-Est du Nigeria. Thème de la biodiversité africaine. Le programme d'Appui à la biodiversité, 1999 ; n°3 pp. 8.
- Okigbo RN, Ogbonnanya OU. Antifungal effects of two tropical plants extracts *Ocimum gratissimum* and *Afromomum melegueta* on post harvest yam *Discorea* spp rot. *Afr. J. Biotechnol.* 2006; 5 (9): 727-731.
- Okoli AS, Okeke MI, Iroegbu CU, Ebo PU. *Antibacterial activity of Harungana madagascariensis leaf extracts. Phytotherapy research*, 2002; 16 :174-179.
- Okunji CO, IWU MM, Jackson JE, Tally JD. Biological activity of saponins from two *Draceana* species. *Adv. Exp. Med. Biol.* 1996; 28: 404-415.
- Okwu DE. Evaluation of chemical composition of medicinal plants belonging to Euphorbiaceae. *Pak. Vet. J.* 2001; 14:160-162.
- Okwute SK. Plant derived pesticidal and antimicrobial agents for use in agriculture. A review of phytochemical and biological studies on some Nigerian plants. *J. Agric. Sci. Technol.* 1992; 2:62-70.
- Omogbai BA, Eneh TO. 2011. Antibacterial activity of *Dacryodes edulis* seed extracts on food-borne pathogens. *Bayero Journal of Pure and Applied Sciences*, 2011; 4(1):17-21.
- Onocha PA, Ekundayo O, Oyelola O, Laakso I. Essential oils of *Dacryodes edulis* (G.Don) H.J. Lam. (African pear). *Flavour Fragrance*, 1999; 14: 135-139.
- Otieno JN, Magadula JJ, Kakudidi E, Kirimhugya C, Orodho J, Okemo P. Use of ethnobotanical criteria for conservation assessment of plants used for respiratory diseases in Lake Victoria Region, Tanzania. *International Journal of Biodiversity and Conservation*, 2011; 3(11): 610-617.
- Otsuki N, Dang NH, Kumagai E, Kondo A, Iwata S, Morimoto C. Aqueous extract of *Carica papaya* leaves exhibits anti-tumor activity and immunomodulatory effects. *Journal of Ethnopharmacology*, 2010; 127: 760–767.
- Oussou KR, Yolou S, Boti JB, Guessennd KN, Kanko C, Ahibo C, Casanova J. Etude chimique et activité anti-diarrhéique des huiles essentielles de deux plantes aromatiques de la pharmacopée ivoirienne. *European Journal of Scientific Research*, 2008 ; 24 (1) : 94-103.
- Owoyale JA, Olatunji GA, Oguntoye SO. Antifungal and antibacterial activities of an alcoholic extract of *Senna alata* leaves. *J. Appl. Sci. Environ. Mgt.* 2005; 9(3):105-107.
- Owoyale JA, Olatunji GA, Oguntoye SO. Antifungal and Antibacterial Activities of an Alcoholic Extract of *Senna alata* Leaves. *J. Appl. Sci. Environ. Mgt.* 2005; 9(3): 105-107.
- Oyewole OI, Massaquoi HO. Oral administration of *Rauwolfia vomitoria* extract has no untoward effect on kidney and liver functions in rats. *African Journal of Biotechnology*, 2008; 7 (10): 1577-1580.
- Panda SK, Thatoi HN, Dutta SK. Antibacterial activity and phytochemical screening of leaf and bark extracts of *Vitex negundo* I. from simimipal biosphere reserve, Orissa. *Journal of Medicinal Plants Research*, 2009; 3(4):294-300.
- Panghal M, Arya, Yadar S, Kumar S, Yadav PJ. Indigenous knowledge of medicinal plants used by Saperas community of khetawas, Jhaffar District, Haryana, India. *Journal of ethnobiology and Ethnomedecine*, 2010; 6: 4.
- Papadimitropoulos V, Vergidis PI, Bliziotis I, Falagas ME. Vaccination against fever in travelers : a cost-effectiveness approach. *Clin. Microbiol. Infect.*, 2004; 10:681-683.

- Parekh J, Chanda SV. Antibacterial activity of aqueous and alcoholic extracts of 34 Indian medicinal plants against some *Staphylococcus* species. *Turk. J. Biol*, 2008; 32: 63-71.
- Parthasarathy R, Ilavarasan R, Karrunakaran CM. Antidiabetic activity of *Thespesia populnea* bark and leaf extract against streptozotocin induced diabetic rats. *Int. J. Pharm. Tech. Res*, 2009; 1(4): 1069-1072.
- Passalacqua NG, Guarrera PM, De Fine G. Contribution to the knowledge of the folk plant medicine in Calabria region (Southern Italy). *Fitoterapia*, 2007; 78:52-68.
- Pettit RK, Weber CA, Kean MJ, Hoffmann H, Pettit GR, Tan R, Franks KS, Horton ML. Microplate alamar blue assay for *Staphylococcus epidermidis* biofilm susceptibility testing. *Antimicrob. Agent Chemother*, 2005; 56: 227-232.
- Pieme CA, Penlap VN, Nkegoum B, Taziebou CL, Tekwu EM, Etoa FX, Ngongang J. Evaluation of acute and subacute toxicities of aqueous ethanolic extract of leaves of *Senna alata* (L.) Roxb (Cesalpiniaceae). *Afr. J. Biotech*, 2006; 5 : 283- 289.
- Pimenta LPS, Mendonça DD, Pretti DL, Cruz BLD, Leite EA, De Oliveira MC. Evaluation of *in vivo* antitumor activity of *Annona crassiflora* wood extract. *International Journal of pharmaceutical Sciences and Drug Research*, 2011; 3(3): 270-273.
- Rabe T, Van Staden J. Antibacterial activity of South African plant used for medicinal purposes. *J. Ethnopharmacol*, 1997; 56: 81-87.
- Ragasa CY, Martinez A, Chua JEY, Rideout JA. A triterpene from *Musa errans*. *Philippine J. Sci*, 2007; 136(2): 167-171.
- Rajesh K, Harsha R, Mohammed GA, Hareesh AR, Thamanna GSS, Dinesha R, Satish KBP, Irfan AM. Antimicrobial activity of ethanol extract of leaf and flower of *Spathodea campanulata* P. Beauv. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2010; 1(3):691-698.
- Rajith NP, Navas M, Thaha AM, Manju MJ, Anish N, Rajasekharan S, George V. A study on traditional mother care plants of rural communities of South Kerala. *Indian Journal of Traditional Knowledge*, 2010; 9(1): 203-208.
- Ravikumar PHS, Makari HK, Gurumurthy H. In vitro antimicrobial activity of ethanol extract of *Thevetia peruviana*. *EJEAFChe*, 2007; 6(9):2318-2322.
- Rindos D.1984. Origins of Agriculture: An evolutionary perspective. Cornell University. Orlando: Academic Press, INC.
- Rojas JJ, Ochoa VJ, Ocampo SA, Munoz JF. Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: A possible alternative in the treatment of non-nosocomial infections. *BMC Complementary and Alternative medicine*, 2006; 6: 2.
- Ronchetti F, Russo G. A new alkaloid from *Rauvolfia vomitoria*. *Phytochemistry*, 1971; 10:1385-1388.
- Rusia K, Srivastava SK. Antimicrobial activity of some Indian medicinal plants. *Ind. J. Pharm. Sci*, 1998; 60:57-58.
- Saravanakumar A, Venkateswaran K, Vanitha J, Ganesh M, Vasudevan M, Sivakumar T. Evaluation of antibacterial activity, phenol and flavonoid contents of *Thespesia populnea* flower extracts. *Pak. J. pharm. Sci*, 2009; 22(3): 282-286.
- Seri-Kouassi BP, Kanko C, Aboua LRN, Bekon KA, Glitho AI, Koukoua G, N'Guessan YT. Action des huiles essentielles de deux plantes aromatiques de Côte d'Ivoire sur *Callosobruchus maculatus* F. du niébé. *C.R. chimie*, 2004 ; 7: 1043-1046.
- Shahid-ud-Daula AFM, Basher MA. Phytochemical screening, plant growth inhibition, and antimicrobial activity studies of *Xylocarpus granatum*. *Malaysian J. Pharm. Sci*, 2009; 7(1): 9-21.
- Siddharth S, Rakesh KS, Sanjeev K, Pradeep K. Anti-psoriatic and phytochemical evaluation of *Thespesia populnea* bark Extracts. *Inter. J. Pharm. and Pharm. Sci*, 2009; 1(1): 176-185.
- Silva FL, Fischer DCH, Tavares JF, Silva MS, De Athayde-Filho PF, Barbosa-Filho JM. Compilation of secondary metabolites from *Bidens pilosa* L. *Molecules*, 2011; 16: 1070-1102.
- Singh R, Singh SK, Arora S. Evaluation of antioxidant potential of ethyl acetate extract/fractions of *Acacia auriculiformis* A. Cunn. *Fod Chem. Toxicol*, 2007; 45: 1216-1223.
- Sofowora A. 1993. Medicinal plants and traditional medicine in Africa. *Spectrum Books Ltd., Ibadan, Nigeria*, pp. 191-289.
- Suffredim IB, Sarder HS, Goncalves AG. 2004. Screening of antibacterial extracts from plants native to Brazilian amazon Rain Forest and Atlantic Forest. *Brazilian J. Med. Biol. Res*, 2004; 37:379-384.
- Suresh K, Saravana Babu S, Harisaranraj R. Studies on in vitro Antimicrobial activity of ethanol extract of *Rauvolfia tetraphylla*. *Ethnobotanical Leaflets*, 2008; 12: 586-90.
- Tadesse M, Hunde D, Getachew Y. Survey of medicinal plants used to treat human diseases in Seka Chekorsa, Jimma Zone, Ethiopia. *Ethiopian Journal of Health Sciences*, 2005; 15(2):89-106. http://ejhs.ju.edu.et/ejhs_journal/2005/volume15issue2.php.
- Tan PV, Dimo T, Dongo E. Effects of Methanol, cyclohexane and methylene chloride extracts of *Bidens pilosa* and various gastric ulcer models in rats. *J. Ethnopharmacol*, 2000; 73: 415-421.
- Taylor JLS, Rabe T, McGraw LJ.2001. Towards the scientific validation of traditional medicinal plants. *Plant Growth Regul*, 2001; 34:23-37.
- Teke NG, Lunga P, Wabo KH, Kuate JR, Vilarem G, Giacinti G, Kikuchi H, Oshima Y. Antimicrobial and antioxidant properties of methanol extract, fractions and compounds from the stem bark of *Entada abyssinica* Stend ex A. Satapie. *BMC Complementary and Alternative Medicine*, 2011; 11:57.
- Tripathi R, Dwivedi SN, Dwivedi S. Ethno-medicinal plants used to treat gynecological disorders by tribal people of Madhya Pradesh, India. *International Journal of Pharmacy & Life Sciences*, 2010; 1(3): 160-169.
- Tsobou R, Mapongmetsem PM, Van Damme P. Medicinal plants used Against typhoid fever in Bamboutos Division, Western Cameroon. *Ethnobotany Research & Applications*, 2013; 11: 163-174.
- Ubillas RP, Mendez CD, Jolad SD, Luo J, King SR, Carlson TJ, Fort DM. Antihyperglycemic acetylenic glucosides from *Bidens pilosa*. *Planta Medica*, 2000; 66: 82-83.
- Ubulom P, Akpabio E, Udobi CE, Mbon R. Antifungal activity of aqueous and ethanolic extracts of *Picralima nitida* seeds on *Aspergillus flavus*, *Candida albicans* and *Microsporum canis*. *Research in Pharmaceutical Biotechnology*, 2011; 3(5): 57-60.
- Venkata ASB, Sai DKS. 2011. Pharmacognostic and phytochemical studies of *Thespesia populnea* Linn. *J. Chem. Pharm. Res*, 2011; 3(4): 237-244.
- Vijayakumar S, Presannakumar G, Vijayalakshmi NR. 2008. Antioxidant activity of banana flavonoids. *Fitoterapia*, 2008; 79: 279-282.
- Wagner KH, Elmadfa I. Biological relevance of terpenoids: Overview focusing on mono-di and tetraterpenes. *Ann Nutr Metab*, 2003; 47:95-106.
- Washington J. 1995. Manual of clinical microbiology. 4th Edn., Lennette, E., A. Balows, W.J. Hausler and H.J. Shadomy (Eds.), American Society of Microbiology, Washington, pp: 967-970.
- World Health Organisation. WHO traditional medicine strategy 2002-2005. World Health Organization, 2002.
- World Health Organization. 2003. Traditional Medicine. [Http://www.who.int/mediacentre/factsheets/fs134/en/print.html](http://www.who.int/mediacentre/factsheets/fs134/en/print.html)
- Wu Y, Hao B, Shen E, Zhang X, Zhao Y. Regulatory effect of *Laggera alata* extract on immune mediated liver injury. *Journal of Medicinal Plants Research*, 2011; 5(12): 2494-2498.
- Yemoa AL, Gbenou JD, Johnson RC, Djego JG, Zinsou C, Moudachirou M, Quetin- Leclercq J, Bigot A, Portael F. 2008. Identification et étude phytochimique des plantes utilisées dans le traitement traditionnel de l'ulcère de Buruli au Bénin. *Ethnopharmacologia*, 2008 ; 42: 48-55.
- Yoga Latha L, Darah I, Sasidharan S, Jain K. Antimicrobial Activity of *Emilia sonchifolia* DC., *Tridax procumbens* L. and *Vernonia cinerea* L. of Asteraceae Family: Potential as food preservatives. *Mal. J. Nutr*, 2009; 15(2): 223-231.

Yusha'a M, Taura, DW, Bello B, Abdullahi N. Screening of *Annona squamosa* extracts for antibacterial activity against some respiratory tract isolates. *International Research of Pharmacy and Pharmacology*, 2011; 1(9):237-241.

Zhao Y. New Eudesmane and Eremophilane derivatives from *Laggera alata*. *Chinese Chemical Letters*, 2003; 14(4): 393-396.

Zofou D, Mofor CT, Zollo PHA. Etude socioépidémiologique du paludisme dans la localité de Mbouda (Ouest-Cameroun). *Medecine d'Afrique Noire*, 2009; 56(2) :91-95.

How to cite this article:

Tsobou Roger, Mapongmetsem Pierre-Marie, Voukeng Kenfack Igor, Van Damme Patrick. Phytochemical screening and antibacterial activity of medicinal plants used to treat typhoid fever in Bamboutos division, West Cameroon. *J App Pharm Sci*, 2015; 5 (06): 034-049.