



Ameliorative effects of *Paederia foetida* Linn. on lead acetate-exposed rats

Tanaporn Khamphaya¹, Phisit Pouyfung¹, Supabhorn Yimthiang¹, Manas Kotepui², Saruda Kuraeiad^{2,3*}

¹Department of Occupational Health and Safety, School of Public Health, Walailak University, Nakhon Si Thammarat, Thailand.

²Department of Medical Technology, School of Allied Health Sciences, Walailak University, Nakhon Si Thammarat, Thailand.

³Research Excellence Center for Innovation and Health Product, Walailak University, Nakhon Si Thammarat, Thailand.

ARTICLE INFO

Received on: 15/08/2021

Accepted on: 12/11/2021

Available Online: 05/03/2022

Key words:

Paederia foetida, lead, hepatoprotective, hematology, liver, kidney.

ABSTRACT

Lead, a toxic heavy metal, induces several health problems via oxidative stress. The *Paederia foetida* extract (PFE), a plant indigenous to Asia, reportedly exhibits potent antioxidant activity. Thereby, this study aimed to investigate the protective effects of PFE on the hematological system and renal and liver function in rats exposed to lead poisoning. Male Wistar rats were divided into six groups and orally administered either 0.25% carboxymethylcellulose as a control, lead acetate [50 mg/kg body weight (BW)], or lead acetate plus one of four concentrations of PFE (50, 100, 500, and 1,000 mg/kg BW). All rats received their treatment daily for 8 weeks. The bodyweight of each animal was recorded weekly. The effects of the lead acetate and PFE were investigated by evaluating the bodyweight and hematological and biochemical parameters, as well as histological studies on the liver and kidney tissues. PFE at 100, 500, and 1,000 mg/kg significantly ($p < 0.05$) improved the harmful effects of lead administration on the bodyweight, the liver, and the kidney, as well as hematological indices. In conclusion, PFE has a potent protective property against lead-induced toxicity. Since oxidative stress results from exposure to lead, an exogenous chelator with antioxidant activity, such as the low-cost PFE, could be a promising agent for ameliorating lead toxicity.

INTRODUCTION

Lead (Pb) is a toxic heavy metal in the environment, and it represents a great concern to human health worldwide (Abdelhamid *et al.*, 2020). It presents in multiple forms and sources, such as food and air pollution (Kosnett *et al.*, 2007). It causes multiorgan toxicities and damages almost all vital organs such as the testes, brain, kidney, liver, and hematopoietic tissues (Manoj Kumar *et al.*, 2017). Lead-contaminated food and water are absorbed through the duodenum via divalent metal transporter 1 and leached into the circulation (Kosnett *et al.*, 2007). More than 95% of the lead in the circulation binds to proteins in the erythrocytes (Gonick, 2011); thus, it is stored in internal organs such as the liver and the kidney (Nakhaee *et al.*, 2019). One of the key mechanisms implicated in lead-induced toxicity is oxidative stress, which is an imbalance between oxidant and antioxidant

systems due to an excess of reactive oxygen species (ROS). This excessive formation of ROS by lead consequently results in mitochondrial impairment and cell damage (Zolkipli-Cunningham and Falk, 2017). Several heavy metal chelating agents have been applied as therapeutic drugs for lead-exposed patients, such as succimer, dimercaprol, dimercaptosuccinic acid, and CaNa₂-ethylenediaminetetraacetic acid (EDTA). However, these chelators have multiple side effects (Kosnett, 2010). Therefore, great attention has been given to natural compounds that could attenuate the deleterious effects of lead poisoning and protect the cell from lead-induced damage. Such natural compounds include vitamin C (Haridy *et al.*, 2014) and vitamin E (Khodamoradi *et al.*, 2015) and flavonoids from natural plants (Adhikari *et al.*, 2018).

Flavonoid and other polyphenol mixtures can be found in several plants, and they are widely known for their antioxidant activity (Pandey and Rizvi, 2009). In a previous experimental animal study, the effects of lead were neutralized by a mixture of flavonoids and polyphenols via heavy-metal chelation and antioxidant activity (Adhikari *et al.*, 2018). Moreover, several flavonoid-containing medicinal plants, such as *Thunbergia laurifolia* Linn. (Tangpong and Satarug, 2010), green tea extract

*Corresponding Author

Saruda Kuraeiad, Department of Medical Technology, School of Allied Health Sciences, Walailak University, Nakhon Si Thammarat, Thailand. E-mail: saruda.ku@mail.wu.ac.th

(Mehana *et al.*, 2012), *Fumaria parviflora* L (Dorostghoal *et al.*, 2014), *Ginkgo biloba* extract (Yallapragada and Velaga, 2015), and *Tinospora cordifolia* (Sharma and Pandey, 2010), also exhibited protective effects against lead-induced toxicity. However, *Paederia foetida* Linn., a flavonoid-enriched plant from the southern region of China, Vietnam, India, Japan, and Thailand (Ahmed *et al.*, 2014) that is used as a folk medicine to treat rheumatoid arthritis (Kumar *et al.*, 2015; Soni *et al.*, 2013), diarrhea (Afroz *et al.*, 2006), asthma (Khushbu *et al.*, 2010), and diabetes (Kumar *et al.*, 2014), has not been investigated for its protective effects on lead-induced toxicity yet. Therefore, this study sought to investigate the protective effects of a methanolic extract of *P. foetida* against low-dose, lead-induced toxicity, particularly in hematologic, renal, and liver organs of rats.

MATERIALS AND METHODS

Ethical approval

The study protocol was approved by the Animal Ethics Committee, Walailak University, under Approval No. 008/2019. All animal procedures were carried out in accordance with the Guide for the Care and Use of Animals for Scientific Purposes, National Committee for Research Animal Development, National Research Council of Thailand.

Paederia foetida Linn. extraction

Paederia foetida leaves were obtained from Khlong Krabue, Pak Phanang District, Nakhon Si Thammarat Province, Thailand (8°17'54.5"N, 100°09'13.1"E); and the leaves were identified by Assistant Professor Dr. Kingkan Bunluepuech, Applied Thai Traditional Medicine Department, School of Medicine, Walailak University, Thailand. The voucher sample (Ref. HA1601061505) was deposited at the School of Medicine, Walailak University. Healthy leaves of *P. foetida* Linn. were collected between March and May 2019. The collected leaves of *P. foetida* were gently washed with tap water. The leaves were dried at 45°C for 48 hours in a hot air oven and then ground into a powder. 50 g of ground leaves was then extracted with 500 ml of methanol for 7 days at room temperature. The plant remains were removed by filtering with Whatman filter paper (No. 1). The filtrated extraction was then concentrated using a rotary vacuum evaporator under reduced pressure at 45°C (Heidolph Hei-VAP Advantage Rotary Evaporator). The concentrated were dried at 45°C in a water bath, resulting in a crude semisolid residue (percentage yield, 20%). The final yield was stored at 4°C for further investigation. In animal studies, it was suspended in a 0.25% solution of carboxymethyl cellulose in distilled water. Doses of the methanolic PFE were modified from Kumar *et al.* (2014, 2018).

Phenolic content of PFE

The amount of phenolic content of PFE was determined using Folin–Ciocalteu's reagent. Briefly, the crude methanolic extract of PFE (1 mg/ml) was mixed with Folin–Ciocalteu's phenol reagent (250: 250 μ l) for 5 minutes. Then, 500 μ l of a sodium carbonate (Na_2CO_3) solution (7% w/v) was added to the mixture,

and the volume was adjusted to 1.5 ml with distilled water. The reactions were allowed to occur for 90 minutes at room temperature (in the dark). Next, 200 μ l of reaction samples was transferred to 96-well microplates, and its absorbance was measured at 765 nm (JASCO FP-8200, JASCO FP-8200 microreader). The amount of phenolic concentration (mg in equivalent/g dry weight) in the plant extract was calculated by using a standard curve that was generated from gallic acid (0–500 μ g/l). All determinations were conducted in triplicate experiments.

Flavonoid content of PFE

To determine the amount of flavonoid content of PFE, 50 μ l of *P. foetida* crude extract (1 mg/ml) was incubated with 200 μ l of distilled water. Then, 5% NaNO_2 (300 μ l) and 10% AlCl_3 (300 μ l) solutions were added, and the final volume was brought to 1 ml. After 5 minutes of incubation, the absorbance was measured at 510 nm and the amount of flavonoid concentration was calculated by using a standard curve that was generated from quercetin (0–500 μ g/l). All determinations were conducted in triplicate experiments.

Lead acetate preparation

Lead (II) acetate trihydrate was obtained from Merck Company, USA. In previous studies, ingestion of low levels of lead at doses of 1, 7.5, 20, 50, and 60 mg/kg bodyweight (BW) caused ROS generation and tissue damage (Agodi *et al.*, 1990; El-Masry *et al.*, 2011; Haridy *et al.*, 2014; Kumar Singh *et al.*, 2018; Offor *et al.*, 2017; Shukla *et al.*, 2003; Tham *et al.*, 2013). Therefore, this study focused on lead acetate at 50 mg/kg BW for lead toxicity induction in rats. A solution was prepared by suspending 5 g of lead acetate in 20 ml of sterile water. A fresh lead acetate solution was prepared daily.

Animals

All animals were obtained from Nomura Siam International, Thailand. They were provided with commercial feed and distilled water. They were maintained in well-ventilated cages, which were kept under a regular light/dark cycle (12/12 hours), constant room temperature (23°C \pm 2°C), and relative humidity (50%–60%). The rats were allowed free access to a standard diet and water throughout the experimental period. The animals were acclimatized for 7 days before the experiments. Bodyweight was recorded weekly for each animal throughout the experiment.

Experimental design

Thirty-six healthy male Wistar rats weighing about 180–200 g at 6 weeks of age were randomly divided into six groups: *i*) animal control group, *ii*) lead-induced group (50 mg/kg BW of lead acetate), *iii*) cotreatment of 50 mg/kg BW PFE with lead acetate group, *iv*) cotreatment of 100 mg/kg BW PFE with lead acetate group, *v*) cotreatment of 500 mg/kg BW PFE with lead acetate group, and *vi*) cotreatment of 1,000 mg/kg BW PFE with lead acetate group.

All rat groups were orally administered selected PFE doses based on previous studies (Arunkumar *et al.*, 2016; Kumar *et al.*, 2014). Treatments lasted for 8 weeks. Eight weeks of lead acetate administration was carried out to mimic subchronic lead

exposure and sufficiency for lead toxicity induction (Obafemi *et al.*, 2019). This also is an appropriate time for investigating the protective effects of PFE (Kumar *et al.*, 2014). The bodyweight of each animal was recorded weekly for 8 weeks.

Dose fixation of the plant drug

The lethal dose or LD₅₀ of the PFE, > 2,000 mg/kg BW, was reported previously (Das *et al.*, 2013; Kumar, 2014). In their study, animals received graded doses of PFE raised from 100 to 2,000 mg/kg BW for 24 hours. They found that no animals showed any change in behavior, anatomy, or neurological patterns from any of the doses. The results from this group suggest that doses up to 2,000 mg/kg of PFE are safe for use. Moreover, several studies used concentrations of PFE between 100 and 500 mg/kg BW for 28 days, and they showed remarkable protective effects, such as antihyperglycemic, antihyperlipidemic, and antioxidant activities, in Wistar rats (Kumar *et al.*, 2014). In addition, at concentrations between 100 and 400 mg/kg BW of PFE, hepatoprotective effects against carbon tetrachloride were seen as well in rats (Arunkumar *et al.*, 2016). Therefore, we chose the dosages of PFE based on these publications and added two more dosages [one lower (50 mg/kg) and one higher (1,000 mg/kg)] to cover the range of promising doses that could protect against lead-induced toxicity.

Blood collection for hematological and biochemical studies

After 8 weeks of treatment, the rats were deeply anesthetized by using a mix of zoletil¹⁰⁰ and xylazine at a ratio of 25:5 mg/kg BW, intraperitoneal injection. The combination of zoletil¹⁰⁰ and xylazine is appropriate to use as an anesthetic since it prevents serum hemolysis and hepatic metabolism in rats (Machado *et al.*, 2009). Blood samples were collected from the heart. Whole blood was collected in EDTA tubes for analyzing complete blood count, including *i*) red blood cell (RBC) count, *ii*) white blood cell (WBC) count, *iii*) hemoglobin concentration (Hb), and *iv*) % hematocrit (Hct), as well as red blood cell indices including *v*) mean corpuscular volume (MCV), *vi*) mean corpuscular hemoglobin (MCH), and *vii*) mean corpuscular hemoglobin concentration (MCHC). These parameters were analyzed by a Mindray Broucher BC-5180 Auto Hematology Analyzer (Aspen Diagnostics, India).

The blood samples were also utilized for biochemical studies. Serum was collected by centrifugation at 3,000 rpm for 30 minutes and then kept in a freezer at -70°C until used. Aspartate aminotransferase (AST) and alanine aminotransferase (ALT), blood urea nitrogen (BUN), and creatinine were evaluated. These parameters were measured by an automated analyzer (Mindray BS-400 Chemistry Analyzer, Aspen Diagnostics, India).

Histopathological study

Histopathological changes in the liver and kidney were determined in this study. The liver and kidney tissues were immediately removed from the rats after blood collection. The tissues were then weighed, fixed in 10% neutral buffered formalin, and embedded in paraffin. After that, sections (5 µm thickness) of paraffin-embedded tissues were stained with hematoxylin and eosin. The histological slides and representative photomicrographs were examined and captured using a light microscope, respectively. The relative weight of the organs was calculated by using the following formula:

$$\text{Relative weight of organ} = \frac{\text{mean absolute weight of organ (g)}}{\text{final weight of the animal (g)}} \times 100$$

Statistical analysis

Data are presented as means ± SEM. Prism 9 software (GraphPad, La Jolla, CA) was used for data analysis. Statistical analyses were carried out using two-tailed unpaired Student's *t*-test when comparing two groups or with a one-way analysis of variance followed by Dunnett's multiple comparisons test when three or more groups were compared. A *p*-value less than 0.05 (*p* < 0.05) was considered statistically significant.

RESULTS

Phenolic and flavonoid content of PFE

According to a previous report, *P. foetida* contains a large amount of flavonoid constituents (Ojha *et al.*, 2018). Therefore, we first determined the phenolic and flavonoid contents in *P. foetida* extracts using garlic and quercetin as the equivalent standard. The phenolic and flavonoid contents of the *P. foetida* extract were 26.47% (3.97 ± 0.17 mg gallic acid equivalents (GAE)/g) and 42.86% (21.43 ± 1.77 mg quercetin equivalents

Table 1. Effects of PFE on bodyweight gain in rats with lead acetate toxicity.

Treatments	Initial BW (g)	Final BW (g)	BW gain (g)	The gain to control (%)
Normal control	225.7 ± 8.4	483.0 ± 16.4	257.3 ± 12.1	100.0 ± 4.7
Lead acetate	220.5 ± 12.0	428.8 ± 18.8	208.3 ± 18.0*	80.9 ± 7.0*
Lead acetate + PFE (50 mg/kg BW)	234.0 ± 6.1	479.3 ± 39.1	245.3 ± 43.0 [#]	95.3 ± 16.7 [#]
Lead acetate + PFE (100 mg/kg BW)	229.8 ± 1.7	473.3 ± 23.7	243.5 ± 24.6 [#]	94.6 ± 9.5 [#]
Lead acetate + PFE (500 mg/kg BW)	231.7 ± 9.3	467.3 ± 32.5	235.7 ± 24.1 [#]	91.6 ± 9.3 [#]
Lead acetate + PFE (1,000 mg/kg BW)	221.0 ± 10.4	457.0 ± 23.1	236.0 ± 13.8 [#]	91.7 ± 5.3 [#]

**p* < 0.05 compared to control group.

[#]*p* < 0.05 compared to lead acetate group.

(QE)/g), respectively, indicating that *P. foetida* contained a good source of flavonoid content.

Effects of lead and PFE on body and organ weight

To determine the toxic effects of PFE on organs and bodyweight, Wistar rats were given lead (50 mg/kg BW) and various concentrations of PFE ranging from 0 to 1,000 mg/kg BW. After 8 weeks of intervention, the result showed that the bodyweight of rats receiving the lead acetate alone exhibited less increase in comparison with rats receiving normal saline (control group) ($p < 0.05$). Moreover, there was no difference in the weight of rats that received lead acetate with any concentration of PFE (50, 100, 500, and 1,000 mg/kg BW) compared to control animals ($p > 0.05$) (Table 1).

The livers and kidneys of the animals taken from the sacrificed animals were weighed and averaged. For the weight of the liver, the results showed no difference in its weight or its weight ratio (weight of liver divided by the bodyweight of rat) in rats exposed to lead acetate compared to controls. Also, there was no difference in the weight or weight ratio of the liver in rats exposed to lead acetate + PFE at any concentration compared to the control group or rats exposed to lead acetate alone. For the weight of the kidney, the results showed no difference in the weight or weight ratio of the kidney in lead-exposed rats compared to controls. However, in Table 2, there was a difference in the weight ratio of the kidney in rats exposed to lead acetate with PFE at 1,000 mg/kg BW compared to controls ($p < 0.05$).

Effects of lead and PFE on hematological parameters

RBC count

The results showed a significant decrease in the RBC count among rats exposed to lead acetate alone compared to controls ($p < 0.05$). Rats exposed to both lead acetate and PFE (at 500 and 1,000 mg/kg BW) exhibited a significant increase in RBC count compared to rats exposed to lead acetate alone ($p < 0.05$).

Hb concentration

The results showed a significant decrease in Hb concentration among rats exposed to lead acetate alone compared to controls ($p < 0.05$). Rats exposed to both lead acetate and PFE (at 50, 100, 500, and 1,000 mg/kg BW) exhibited a significant

increase in Hb concentration compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 1B).

Hematocrit

The results showed a significant decrease in Hct among rats exposed to lead acetate alone compared to controls ($p < 0.05$). Rats exposed to both lead acetate and PFE (at 50, 100, 500, and 1,000 mg/kg BW) exhibited a significant increase in Hct compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 1C).

Mean corpuscular volume

The results showed a significant decrease in MCV among rats exposed to lead acetate alone compared to those in the control ($p < 0.05$). Rats exposed to both lead acetate and PFE (at 500 and 1,000 mg/kg BW) exhibited a significant increase in MCV compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 1D).

Mean corpuscular hemoglobin

The results showed a significant decrease in MCH among rats exposed to lead acetate alone compared to controls ($p < 0.05$). There was no difference in MCH between rats exposed to both lead acetate and PFE (at 50, 100, 500, and 1,000 mg/kg BW) and rats exposed to lead acetate alone ($p > 0.05$) (Fig. 1E).

Mean corpuscular hemoglobin concentration

The results showed a significant decrease in MCHC among rats exposed to lead acetate alone compared to those in the control ($p < 0.05$). Rats exposed to both lead acetate and PFE (at 100, 500, and 1,000 mg/kg BW) exhibited a significant increase in MCHC compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 1F).

Red cell distribution width (RDW)

The results showed no difference in RDW between rats exposed to lead acetate alone and rats in the control ($p > 0.05$). There was also no difference in MCH between rats exposed to both lead acetate and PFE (at 50, 100, 500, and 1,000 mg/kg BW) and rats exposed to lead acetate alone ($p > 0.05$) (Fig. 1G).

WBC Count

The results showed a significant increase in WBC count among rats exposed to lead acetate alone compared to controls ($p < 0.05$). However, rats exposed to both lead acetate and PFE

Table 2. Effects of PFE on organ weight ratio in rats induced with lead acetate.

Treatments	BW	Liver			Kidney		
		Weight (g)	Ratio (%)	%	Weight (g)	Ratio (%)	%
Normal control	483.0 ± 16.4	15.4 ± 0.6	3.2 ± 0.1	100	3.2 ± 0.4	0.66 ± 0.06	100
Lead acetate	428.8 ± 18.8	14.87 ± 1.2	3.5 ± 0.2	109	3.4 ± 0.4	0.78 ± 0.06	118
Lead acetate + PFE (50 mg/kg BW)	479.3 ± 39.1	15.3 ± 1.9	3.3 ± 0.3	104	3.4 ± 0.3	0.76 ± 0.05	115
Lead acetate + PFE (100 mg/kg BW)	473.3 ± 23.7	17.8 ± 3.6	3.7 ± 0.6	115	3.5 ± 0.3	0.74 ± 0.09	112
Lead acetate + PFE (500 mg/kg BW)	467.3 ± 32.5	14.9 ± 0.8	3.2 ± 0.1	98	3.5 ± 0.5	0.76 ± 0.14	115
Lead acetate + PFE (1,000 mg/kg BW)	457.0 ± 23.1	14.5 ± 0.5	3.2 ± 0.6	99	3.8 ± 0.3	0.83 ± 0.04*	125*

* $p < 0.05$ compared to control group.




Figure 1. Effects of lead acetate and the PFE on hematological parameters. (A) Red blood cell (RBC) count; (B) Hemoglobin (Hb) concentration; (C) % hematocrit (Hct); (D) mean corpuscular volume (MCV); (E) mean corpuscular hemoglobin (MCH); (F) mean corpuscular hemoglobin concentration (MCHC); (G) % red cell distribution width (RDW); (H) white blood cell (WBC) count; PFE: methanol extract of *Paederia foetida* leaves (mg/kg BW); lead (50 mg/kg BW). * $p < 0.05$ compared to control, # $p < 0.05$ compared between the lead acetate and PFE cotreatment groups.

(at 50, 100, 500, and 1,000 mg/kg BW) exhibited a significant decrease in WBC count compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 1H).

Effects of PFE on liver function and histopathology

Alanine aminotransferase

The results showed that rats exposed to lead acetate alone exhibited significantly increased ALT compared to controls

($p < 0.05$) (Fig. 2A). However, rats exposed to lead acetate with PFE (at 50, 100, 500, and 1,000 mg/kg BW) exhibited significantly decreased ALT compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 2A).

Aspartate aminotransferase

The results showed that rats exposed to lead acetate alone exhibited significantly increased AST compared to controls




Figure 2. Effects of the PFE on liver function and histopathology. (A) Serum aspartate aminotransferase (AST) level; (B) serum alanine aminotransferase (ALT) level. * $p < 0.05$ compared to the control, # $p < 0.05$ compared between the lead acetate and the PFE cotreatment groups. Photographs show representative hepatocytes stained with hematoxylin and eosin. (C) Hepatocytes in the control group; (D) hepatocytes in the lead acetate group; (E) hepatocytes in lead + PFE 50 mg/kg BW group; (F) hepatocytes in lead + PFE 100 mg/kg BW group; (G) hepatocytes in lead + PFE 500 mg/kg BW group; and (H) hepatocytes in lead + PFE 1,000 mg/kg BW group. Scale bar = 40 μ m (magnification = 400 \times).

($p < 0.05$) (Fig. 2B). Rats exposed to lead acetate with PFE (at 100, 500, and 1,000 mg/kg BW) exhibited significantly decreased AST compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 2B).

Histopathology

No histopathological abnormalities were observed in the hepatocytes of rats in the control group (Fig. 2C). Rats exposed to lead acetate alone showed some neutrophil infiltration around the periportal areas (Fig. 2D). However, no histopathological abnormalities were observed in the hepatocytes of rats exposed to lead acetate and PFE at 50 mg/kg BW (Fig. 2E), 100 mg/kg BW (Fig. 2F), 500 mg/kg BW (Fig. 2G), and 1,000 mg/kg BW (Fig. 2H).

Effects of PFE on kidney function and histopathology

Blood urea nitrogen

The results showed that rats exposed to lead acetate alone exhibited significantly increased BUN compared to the control group ($p < 0.05$). Rats exposed to lead acetate with PFE (at 100, 500, and 1,000 mg/kg BW) exhibited significantly decreased BUN compared to rats exposed to lead acetate alone ($p < 0.05$) (Fig. 3A).

Creatinine

The results showed no difference in creatinine levels between rats exposed to lead acetate alone and the control group ($p > 0.05$). Also, there was no difference in creatinine levels between rats exposed to both lead acetate and PFE (at 50, 100, 500, and 1,000 mg/kg BW) and rats exposed to lead acetate alone ($p > 0.05$) (Fig. 3B).

Kidney histopathology

No histopathological abnormalities could be observed in the kidneys of rats in the control group (Fig. 3C). However, there was mild parenchymal cell degeneration among rats exposed to lead acetate alone (Fig. 3D) and rats exposed to both lead acetate and PFE at 50 mg/kg BW (Fig. 3E), 100 mg/kg BW (Fig. 3F), 500 mg/kg BW (Fig. 3G), and 1,000 mg/kg BW (Fig. 3H).

DISCUSSION

Lead is a highly toxic heavy metal that has a variety of pathological and physiological effects in both animals and humans (El-Boshy *et al.*, 2019; El-Tantawy, 2016). In this work, lead acetate-induced hepato-, renal- and hematotoxicities in rats. However, the toxic and harmful effects of lead acetate were alleviated by the *P. foetida* extract (PFE) starting at the dose of 100 mg/kg BW, which may have a higher potency and similar efficacy when compared with higher doses. PFE at a dose of 100 mg/kg BW also showed several other effective characteristics; for example, *i*) administration of PFE for 5 days significantly prevented hepatotoxicity after injection of CCl_4 for 48 hours in rats (Arunkumar *et al.*, 2016) and *ii*) PFE at a daily dose of 100 mg/kg for 28 days significantly decreased the level of fasting blood glucose, total cholesterol, and triglyceride, and it modulated the pathology of the kidney and liver in streptozotocin-induced diabetic rats (Kumar *et al.*, 2014). Moreover, the maximum

effective dose observed in diabetic rats was 500 mg/kg BW, which was slightly aligned with our hematology results. It has also been reported that PFE can protect against colitis in rats at the higher dose of 500 mg/kg BW (Das *et al.*, 2013). It seems that the higher concentration of PFE was also safe and effective and showed no signs of toxicity as observed in our studies and previous reports, which found that PFE doses up to 2,000 mg/kg BW (p.o.) are safe with an LD_{50} of more than 2,000 mg/kg BW (Das *et al.*, 2013; Kumar *et al.*, 2014).

The present study proposed that PFE could attenuate the lead-induced reduction in bodyweight and restore hematologic alteration, renal function, and liver function in lead acetate-induced rats. For lead-induced reductions in bodyweight, the data from National Health and Nutrition Examination Survey (NHANES) 1999–2006 revealed that blood lead levels were associated with lower bodyweights in adults (Scinicariello *et al.*, 2013), which can potentially be caused by lead-induced metabolic changes. Moreover, previous studies indicated subchronic exposure to heavy metals induced weight loss (Nwokocha *et al.*, 2011, 2012). The mechanism of lead-induced weight loss is not well understood. A previous study suggested that the loss of bodyweight might be caused by the lead-induced oxidative stress in the muscular system through ROS generation, leading to muscular atrophy (Buck and Chojkier, 1996). Following PFE treatment in lead-exposed rats, the present study showed that PFE in any concentration could improve the bodyweight of lead-exposed rats when compared to lead-exposed rats without PFE. The antioxidant properties of PFE could potentially explain its weight-restorative effect among lead-exposed rats.

The present study also showed no difference in the weight or weight ratio of the internal organs (liver and kidney) between rats exposed to lead acetate alone, the control group, or rats exposed to lead acetate + PFE. These results are partially consistent with the study of Amjad *et al.*, (2013), who reported that the weight of the kidney was significantly increased at 4 weeks, following daily induction (i.p.) of lead acetate at 8 mg/kg BW, and it did not change significantly at 6 weeks. Therefore, the alteration of the internal organs among lead-exposed rats is dependent on the duration of exposure. The mechanism underlying the increased internal organ weight could be explained by lipid accumulation in the liver and kidney, resulting in an increased weight of those organs (Alwaleedi, 2016).

For liver function and histopathology, the present study showed that AST and ALT levels were significantly increased in lead-exposed rats compared to control, suggesting lead-mediated hepatic damage. Moreover, neutrophil infiltration could be observed around the periportal areas of liver tissue in lead-exposed rats. Interestingly, PFE administration at 100, 500, and 1,000 mg/kg BW decreased ALT and AST and reversed or prevented the histopathology of the liver tissue. These results suggest that PFE can protect against hepatic injuries and prevent the hepatic alterations that are caused by the toxic effects of lead acetate exposure. These results are consistent with the findings of previous studies that demonstrated the hepatoprotective activity of PFE in hepatotoxin-induced rats (Arunkumar *et al.*, 2016; Sumithra *et al.*, 2014; Uddin *et al.*, 2011). The mechanism of lead-induced hepatic injuries is not well understood. Previous studies showed that lead acetate could induce the production of ROS-




Figure 3. Effects of PFE on kidney function and histopathology. (A) Blood urea nitrogen (BUN) level; (B) serum creatinine level. * $p < 0.05$ compared to the control, # $p < 0.05$ compared between the lead acetate and the PFE cotreatment groups. Photographs show representative renal cells stained with hematoxylin and eosin. (C) Renal cells control group; (D) renal cells in the lead acetate group; (E) renal cells in lead + PFE 50 mg/kg BW group; (F) renal cells in lead + PFE 100 mg/kg BW group; (G) renal cells in lead + PFE 500 mg/kg BW group; (E) renal cells in lead + PFE 1,000 mg/kg BW group. Scale bar = 40 μ m (magnification = 400 \times). \blacktriangle = parenchymal cell degeneration.

related oxidative damage to the cell membrane of hepatocytes, which leads to hepatic enzyme leakage (Poli *et al.*, 1987; Wang *et al.*, 2013).

For the kidney function and histopathology, the present study showed an increased BUN but not creatinine levels among the rats exposed to lead acetate alone. These results indicate that the alteration in BUN was more sensitive than those of creatinine for the early stages of the lead-induced renal impairment. This was corroborated by another study that showed that increased creatinine was frequent in moderate to severe renal dysfunction (Wong Vega *et al.*, 2020). In addition, mild parenchymal cell degeneration was observed among rats exposed to lead acetate alone (Fig. 3D). Following the administration of lead + PFE at any dose, BUN was significantly decreased (Fig. 3A). Interestingly, mild parenchymal cell degeneration was also observed among rats exposed to lead acetate alone or both lead acetate and PFE. As BUN and creatinine are commonly used as renal impairment biomarkers, the increased BUN level might be an early sign of renal dysfunction. Decreased BUN after lead acetate administration suggests the possibility of the protective effect of PFE.

To determine the long-term effects of lead plus *P. foetida* intake in humans, low-dose lead acetate with *P. foetida* was administered for 8 weeks rats. In this study, lead acetate at a dose of 50 mg/kg BW caused mild liver injury, which is slightly different from other groups that showed more liver damage in rats (Abdelhamid *et al.*, 2020; Adhikari *et al.*, 2018; Offor *et al.*, 2017; Sumithra *et al.*, 2014). This discrepancy could be due to the shorter time point at which the other groups observed the effect of lead exposure (4 weeks). In addition, another report found that 2% lead acetate, which is a subtoxic concentration, causes changes in hepatocytes, portal triads, and sinusoids with an aggregation of inflammatory cells in the portal triad one month after lead exposure (Jarrar and Taib, 2012). However, the liver normally provides a great ability to tolerate several kinds of injuries and toxicants. After chemically induced liver damage, the process of inflammation can neutralize injurious agents and restore liver function. Therefore, the liver might have been repaired and recovered when exposed to lead for a long time, as seen by our liver histology slides that showed only mild liver injury (Fig. 2). This result aligns with another report of normal portal hepatic tissue with a mild aggregation of inflammatory cells in the liver in rats seven months after lead exposure (Jarrar and Taib, 2012).

In the hematological study, the results showed a significant decrease in the hematological parameters, including the RBC count, Hb concentration, % hematocrit, MCV, and MCHC in the lead-exposed rats compared to controls. The results of this study were consistent with the findings of previous studies (Haridy *et al.*, 2014; Obafemi *et al.*, 2019). However, PFE cotreatment with lead acetate showed effective improvement in these hematological parameters. The mechanisms of lead-induced hematological alterations in rats are well described, including *i*) lead-related RBC membrane fluidity that leads to increased erythrocyte hemolysis rates (Mannem, 2014; Ray, 2016), *ii*) lead interference with the heme synthesis pathway, such as the downregulation of δ -aminolevulinic acid dehydratase, resulting in low Hb concentration (Obafemi *et al.*, 2019), or *iii*) the interference of lead with iron and copper metabolism in the RBCs

(Klauder and Petering, 1977). While most of the RBC parameters were altered, the present study showed that the leukocyte count was also increased in lead-exposed rats. This result is consistent with another study, which also showed a lead-induced increase in leukocyte count (Shah and Altindog, 2005). Following PFE administration at any concentration, our results showed restoration of leukocyte counts back to the physiological levels. This result suggests that PFE can reduce the inflammatory responses that are induced by lead toxicity.

To the best of our knowledge, this study is the first report of PFE effectiveness against lead-induced toxicity. A possible mechanism could be due to the effect of various phytochemicals found within PFE (Ojha *et al.*, 2018; Wang *et al.*, 2014). For example, flavonoids such as kaempferol (Satapathy and Pattnaik, 2019) found in PFE could retain their heavy-metal binding activity and act as chelating agents that neutralize the Pb^{2+} charge (Adhikari *et al.*, 2018; Bagchi *et al.*, 2015; Cornard and Merlin, 2003). This fact could reduce ROS and other free-radical-induced oxidative stress and inflammation during Pb^{2+} exposure in the cells. Flavonoids also retain their antioxidant and anti-inflammation properties after Pb^{2+} chelation resulting in even more health benefits. Since significant oxidative stress is generated due to cellular Pb^{2+} exposure, an exogenously supplied chelator with sustainable antioxidant activity and a low cost, like PFE, could be a promising agent for treating lead-induced diseases. Moreover, oxidative stress markers, such as protein carbonylation and nonprotein thiols, and antioxidant enzymes, such as catalase and superoxide dismutase, need to be further evaluated in *P. foetida*. Finally, future studies should include clinical trials to assess the protective effects of *P. foetida* against high blood lead levels in lead-exposed workers.

CONCLUSION

Our investigation clearly indicated that methanolic *P. foetida* leaf extracts, at doses of 100, 500, and 1,000 mg/kg BW, improve bodyweight and the function of the liver, renal, and hematologic systems, following low-level exposure to lead. This study is the first to demonstrate the effects of methanolic extracts from *P. foetida* leaves on the hematologic system and the function of the liver and kidney in lead acetate-induced rats. PFE starting at the doses of 100 mg/kg BW might be beneficial against low doses of lead exposure in rats. However, the potential protective effects of dietary supplements containing PFE following lead poisoning in humans need to be investigated further.

ACKNOWLEDGMENTS

The authors would like to express their sincere gratitude to Associate Professor Dr. Jitbanjong Tangpong and Associate Professor Dr. Wiyada Kwanhian Klangbud for providing some laboratory equipment.

CONFLICTS OF INTEREST

The authors report no financial or any other conflicts of interest related to this work.

FUNDING

This work was supported by the Research Institute for Health Sciences, Walailak University under Grant [number WU-IRG-62-011].

AUTHORS' CONTRIBUTIONS

T.K., P.P., S.Y., and S.K. carried out the experiments; T.K., P.P., and S.K. analyzed the data; T.K., P.P., S.Y., M.K., and S.K. interpreted the results of the experiments. T.K. and S.K. prepared the figures; T.K., P.P., S.Y., M.K., and S.K. drafted, edited, revised, and approved the final version of the manuscript; T.K., P.P., S.Y., and S.K. conceived and designed the research.

ETHICAL APPROVAL

The study protocol was approved by the Animal Ethics Committee, Walailak University (Protocol Number: 008/2019).

REFERENCES

- Abdelhamid FM, Mahgoub HA, Ateya AI. Ameliorative effect of curcumin against lead acetate-induced hemato-biochemical alterations, hepatotoxicity, and testicular oxidative damage in rats. *Environ Sci Pollut Res*, 2020; 27:10950–65.
- Adhikari A, Darbar S, Chatterjee T, Das M, Polley N, Bhattacharyya M, Bhattacharya S, Pal D, Pal SK. Spectroscopic studies on dual role of natural flavonoids in detoxification of lead poisoning: bench-to bedside preclinical trial. *ACS Omega*, 2018; 3(11):15975–87.
- Afroz S, Alamgir M, Khan MT, Jabbar S, Nahar N, Choudhuri MS. Antidiarrhoeal activity of the ethanol extract of *Paederia foetida* Linn. (Rubiaceae). *J Ethnopharmacol*, 2006; 105(1–2):125–30.
- Agodi A, Viola M, Alberghina M, Giaffri da Stella AM. Effect of low-dose lead acetate exposure on the metabolism of nucleic acids and lipids in cerebellum and hippocampus of rat during postnatal development. *J Neurosci Res*, 1990; 25(1):131–8.
- Ahmed A, Islam M, Rahman A, Hossain A. Thrombolytic, cytotoxic and antidiabetic effects of *Paederia foetida* L. leaf extract. *J Adv Med Res*, 2014; 4(5):1244–56.
- Alwaleedi S. Hematobiochemical changes induced by lead intoxication in male and female albino mice. *Natl J Physiol Pharm Pharmacol*, 2016; 6(1):46–51.
- Amjad Z, Iqbal M, Shoro A. Lead-induced reduction in body and kidney weight of Wistar albinorats ameliorated by *Ginkgo biloba* extract (EGb 761). *Biochem Physiol*, 2013; 2(2):1–4.
- Arunkumar R, Chowdhury A, Chowdhury B. Hepatoprotective activity of *Paederia foetida* *in vitro* and *in vivo* studies. *J Evid Based Med Healthc*, 2016; 3(18):733–7.
- Bagchi D, Chaudhuri S, Sardar S, Choudhuri S, Polley N, Lemmens P, Pal SK. Modulation of stability and functionality of a phyto-antioxidant by weakly interacting metal ions: curcumin in aqueous solution. *RSC Adv*, 2015; 5(124):102516–24.
- Buck M, Chojkier M. Muscle wasting and dedifferentiation induced by oxidative stress in a murine model of cachexia is prevented by inhibitors of nitric oxide synthesis and antioxidants. *EMBO J*, 1996; 15(8):1753–65.
- Cornard JP, Merlin JC. Comparison of the chelating power of hydroxyflavones. *J Mol Struct*, 2003; 651:381–7.
- Das S, Kanodia L, Mukherjee A, Hakim A. Effect of ethanolic extract of leaves of *Paederia foetida* Linn. on acetic acid induced colitis in albino rats. *Indian J Pharmacol*, 2013; 45(5):453–7.
- Dorostghoal M, Seyyednejad SM, Jabari A. Protective effects of *Fumaria parviflora* L. on lead-induced testicular toxicity in male rats. *Andrologia*, 2014; 46(4):437–46.
- El-Boshy ME, Refaat B, Qasem AH, Khan A, Ghaith M, Almasmoum H, Mahbub A, Almainani RA. The remedial effect of *Thymus vulgaris* extract against lead toxicity-induced oxidative stress, hepatorenal damage, immunosuppression, and hematological disorders in rats. *Environ Sci Pollut Res Int*, 2019; 26(22):22736–46.
- El-Masry TA, Emar AM, El-Shitany NA. Possible protective effect of propolis against lead-induced neurotoxicity in animal model. *J Evol Biol Res*, 2011; 3(1): 4–11.
- El-Tantawy WH. Antioxidant effects of Spirulina supplement against lead acetate-induced hepatic injury in rats. *J Tradit Complement Med*, 2016; 6(4):327–31.
- Gonick HC. Lead-binding proteins: a review. *J Toxicol*, 2011; 2011:686050.
- Haridy M, Al-Amgad Z, Sakai H, Mohi-Eldin M. Ameliorating effects of garlic, calcium, and vitamin C on chronic lead toxicity in albino rats. *Comp Clin Pathol*, 2014; 23(5):1215–23.
- Jarrar BM, Taib NT. Histological and histochemical alterations in the liver induced by lead chronic toxicity. *Saudi J Biol Sci*, 2012; 19(2):203–10.
- Khodamoradi N, Komaki A, Salehi I, Shahidi S, Sarihi A. Effect of vitamin E on lead exposure-induced learning and memory impairment in rats. *Physiol Behav*, 2015; 144:90–4.
- Khushbu C, Anar P, Mayuree P, Carol M, Roshni S, Subodh A. *Paederia foetida* Linn. as a potential medicinal plant: a review. *J Pharm Res*, 2010; 3(12):3135–7.
- Klauder DS, Petering HG. Anemia of lead intoxication: a role for copper. *J Nutr*, 1977; 107(10):1779–85.
- Kosnett MJ. Chelation for heavy metals (arsenic, lead, and mercury): protective or perilous? *Clin Pharmacol Ther*, 2010; 88(3):412–5.
- Kosnett MJ, Wedeen RP, Rothenberg SJ, Hipkins KL, Materna BL, Schwartz BS, Hu H, Woolf A. Recommendations for medical management of adult lead exposure. *Environ Health Perspect*, 2007; 115(3):463–71.
- Kumar Singh P, Kumar Singh M, Singh Yadav R, Kumar Dixit R, Mehrotra A, Nath R. Attenuation of lead-induced neurotoxicity by Omega-3 fatty acid in rats. *Ann Neurosci*, 2018; 24(4):221–32.
- Kumar V, Al-Abbasi FA, Ahmed D, Verma A, Mujeeb M, Anwar F. *Paederia foetida* Linn. inhibits adjuvant induced arthritis by suppression of PGE(2) and COX-2 expression via nuclear factor-kappaB. *Food Funct*, 2015; 6(5):1652–66.
- Kumar V, Anwar F, Ahmed D, Verma A, Ahmed A, Damanhoury ZA, Mishra V, Ramteke PW, Bhatt PC, Mujeeb M. *Paederia foetida* Linn. leaf extract: an antihyperlipidemic, antihyperglycaemic and antioxidant activity. *BMC Complement Altern Med*, 2014; 14:76.
- Kumar V, Kaithwas G, Anwar F, Rahman M, Patel DK, Singh Y, Verma A. Effect of variable doses of *Paederia foetida* L. combat against experimentally-induced systemic and topical inflammation in Wistar rats. *Curr Bioact Compd*, 2018; 14(1):70–9.
- Machado EF, Normand AC, Nunes LA, Brenzikof R, Macedo DV. Effects of different general anesthetics on serum hemolysis and hepatic and muscular glycogenolysis in rats. *Braz J Med Biol Res*, 2009; 42(11):1035–8.
- Mannem P. Lead toxicity on hematological changes and amelioration with ginger (*Zingiber officinale*) extract in male albino rats. *Int J Adv Res*, 2014; 2(4):23–8.
- Manoj Kumar V, Henley AK, Nelson CJ, Indumati O, Prabhakara Rao Y, Rajanna S, Rajanna B. Protective effect of *Allium sativum* (garlic) aqueous extract against lead-induced oxidative stress in the rat brain, liver, and kidney. *Environ Sci Pollut Res*, 2017; 24(2):1544–52.
- Mehana EE, Meki AR, Fazili KM. Ameliorated effects of green tea extract on lead induced liver toxicity in rats. *Exp Toxicol Pathol*, 2012; 64(4):291–5.
- Nakhaee S, Amirabadizadeh A, Brent J, Mehrpour O. Impact of chronic lead exposure on liver and kidney function and hematologic parameters. *Basic Clin Pharmacol Toxicol*, 2019; 124(5):621–8.
- Nwokocho CR, Nwokocho MI, Owu DU, Obi J, Olatunde B, Ebe C, Nwangwu O, Iwuala MO. Comparative analysis on the effect of palm oil (*Elaeis guineensis*) in reducing cadmium and lead accumulation in liver of Wistar rats. *Pharmacognosy Res*, 2012; 4(4):214–8.

- Nwokocha CR, Owu DU, Ufearo CS, Iwuala MOE. Comparative study on the efficacy of *Garcinia kola* in reducing some heavy metal accumulation in liver of Wistar rats. *J Ethnopharmacol*, 2011; 135(2):488–91.
- Obafemi TO, Onasanya A, Adeoye A, Falode JA, Daniel DJ, Irefo EF, Ojo AO, Fadaka A, Afolabi OB, Awe JO, Omiyale BO. Protective effect of methanolic and flavonoid-rich leaf extracts of *Synsepalum dulcificum* (Danielli) on lead-acetate-induced toxicity in Wistar albino rats. *J Appl Pharm Sci*, 2019; 9(5):65–72.
- Offor SJ, Mbagwu HO, Orisakwe OE. Lead induced hepatorenal damage in male albino rats and effects of activated charcoal. *Front Pharmacol*, 2017; 8:107.
- Ojha S, Raj A, Roy A, Roy S. Extraction of total phenolics, flavonoids and tannins from *Paederia foetida* L. leaves and their relation with antioxidant activity. *Pharmacogn J*, 2018; 10(3):541–7.
- Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev*, 2009; 2(5):270–8.
- Poli G, Albano E, Dianzani MU. The role of lipid peroxidation in liver damage. *Chem Phys Lipids*, 1987; 45(2–4):117–42.
- Ray RR. Haemotoxic effect of lead: a review. *Proc Zoo Soc*, 2016; 69(2):161–72.
- Satapathy S, Pattnaik G. Antioxidant and HPLC analysis of an Indian medicinal herb: *Paederia foetida* L. (Prasarini). *Int J Life Sci*, 2019; 7(2):249–55.
- Scinicariello F, Buser MC, Mevissen M, Portier CJ. Blood lead level association with lower body weight in NHANES 1999–2006. *Toxicol Appl Pharmacol*, 2013; 273(3):516–23.
- Shah SL, Altindog A. Alteration in the immunological parameters Tench (*Tinca tinca* L. 1758) after acute and chronic exposure to lethal and sublethal treatments with mercury, cadmium and lead. *Turk J Vet Anim Sci*, 2005; 29:1163–8.
- Sharma V, Pandey D. Protective role of *Tinospora cordifolia* against lead-induced hepatotoxicity. *Toxicol Int*, 2010; 17(1):12–7.
- Shukla PK, Khanna VK, Khan MY, Srimal RC. Protective effect of curcumin against lead neurotoxicity in rat. *Hum Exp Toxicol*, 2003; 22(12):653–8.
- Soni RK, Irchhaiya R, Dixit V, Alok S. *Paederia foetida* Linn: phytochemistry, pharmacological and traditional uses. *Int J Pharm Sci Res*, 2013; 4(12):4525.
- Sumithra M, Chitra V, Anbu J, Suman R, Shubaranshu, Nithya S. Hepatoprotective activity of leaf extract of *Paederia foetida* in experimental liver cirrhosis. *Invent Rapid Ethnopharmacol*, 2014; (2):1–6.
- Tangpong J, Satarug S. Alleviation of lead poisoning in the brain with aqueous leaf extract of the *Thunbergia laurifolia* (Linn.). *Toxicol Lett*, 2010; 198(1):83–8.
- Tham CS, Chakravarthi S, Haleagrahara N, De Alwis R. Morphological study of bone marrow to assess the effects of lead acetate on haemopoiesis and aplasia and the ameliorating role of *Carica papaya* extract. *Exp Ther Med*, 2013; 5(2):648–52.
- Uddin B, Nahar T, Basunia MA, Hossain S. *Paederia Foetida* protects liver against hepatotoxin-induced oxidative damage. *Adv Biol Res*, 2011; 5(5):267–72.
- Wang J, Zhu H, Yang Z, Liu Z. Antioxidative effects of hesperetin against lead acetate-induced oxidative stress in rats. *Indian J Pharmacol*, 2013; 45(4):395–8.
- Wang L, Jiang Y, Han T, Zheng C, Qin L. A phytochemical, pharmacological and clinical profile of *Paederia foetida* and *P. scandens*. *Nat Prod Commun*, 2014; 9(6):879–86.
- Wong Vega M, Swartz SJ, Devaraj S, Poyyapakkam S. Elevated serum creatinine: but is it renal failure? *Pediatrics*, 2020; 146(1):e20192828.
- Yallapragada PR, Velaga MK. Effect of *Ginkgo biloba* Extract on lead-induced oxidative stress in different regions of rat brain. *J Environ Pathol Toxicol Oncol*, 2015; 34(2):161–73.
- Zolkipli-Cunningham Z, Falk MJ. Clinical effects of chemical exposures on mitochondrial function. *Toxicology*, 2017; 391:90–9.

How to cite this article:

Khamphaya T, Pouyfung P, Yimthiang S, Kotepui M, Kuraeiad S. Ameliorative effects of *Paederia foetida* Linn. on lead acetate-exposed rats. *J Appl Pharm Sci*, 2022; 12(03):160–170.