Journal of Applied Pharmaceutical Science Vol. 14(08), pp 078-084, August, 2024 Available online at http://www.japsonline.com DOI: 10.7324/JAPS.2024.175441 ISSN 2231-3354



Suppression of inflammation by *Baccharis punctulata and Baccharis trimera* through modulation of innate and adaptive immune responses

Christa Burgos¹, Nelson Alvarenga², Pablo H. Sotelo¹, Patricia Langjahr^{1*}

¹Departmento de Biotecnología, Facultad de Ciencias Químicas, Universidad Nacional de Asunción, San Lorenzo, Paraguay. ²Departmento de Fitoquímica, Facultad de Ciencias Químicas, Universidad Nacional de Asunción, San Lorenzo, Paraguay.

ARTICLE HISTORY

Received on: 02/03/2024 Accepted on: 22/06/2024 Available Online: 05/08/2024

Key words:

Baccharis, Asteraceae, inflammation, innate immune response, adaptive immune response, anti-inflammatory agents, medicinal plants.

ABSTRACT

Inflammatory processes, involving both innate and adaptive responses, are essential for controlling pathogens and maintaining homeostasis. However, excessive inflammation can cause chronic conditions including autoimmune diseases, atherosclerosis, and cancer. Alternative treatment options for inflammation are crucial, and exploring plants with anti-inflammatory properties holds significance. The genus *Baccharis* (Asteraceae family) is widespread across the Americas and is traditionally used to treat various disorders, including inflammatory diseases. Several biological activities of *Baccharis* species have been described; however, their immunomodulatory effects have not been widely evaluated. In this study, we analyzed the immunomodulatory activity of two *Baccharis* species, *Baccharis punctulata*, and *Baccharis trimera*, using an *in vitro* inflammation model involving monocytic cells and splenocytes to evaluate both innate and adaptive immune responses. Both *B. punctulata* and *B. trimera* reduced lipopolysaccharide (LPS)-induced inflammatory mediators, including soluble CD14 and pro-inflammatory cytokines, in THP-1 cells. Furthermore, they suppressed nitric oxide (NO) production in splenocytes, demonstrating a dampened innate immune response. In addition, both species attenuated concanavalin A (ConA)-induced splenocyte proliferation, suggesting an anti-inflammatory effect on the adaptive immune response. In summary, the extracts demonstrated significant anti-inflammatory activity, affecting both innate and adaptive immune responses, and underscoring their potential as treatments for inflammatory diseases and sources of anti-inflammatory molecules.

INTRODUCTION

A diverse array of plants has been employed in traditional medicine since ancient times, with a substantial proportion specifically harnessed for the alleviation of inflammatory conditions such as rheumatic diseases and diabetes [1]. For example, in South America, *Baccharis. trimera*, *Annona neosalicifolia*, and *Artemisia absinthium* L. CTES0030264 are used frequently for the treatment of diabetes and *Dysphania ambrosioides*, *Sida cordifolia*, and *Maytenus ilicifolia* as anti-inflammatory agents [2].

Inflammation is a protective reaction that contributes to the return to a homeostatic state after injury. Innate and adaptive immune cells participate in this response [3]. In this context, monocytes/macrophages play a significant role in inflammation, defense mechanisms, and removal of damaged cells. Excessive responses of these cells, on the other hand, can lead to chronic inflammatory conditions including autoimmune diseases, atherosclerosis, and cancer, among others. Overproduction of inflammatory mediators by monocytes, such as pro-inflammatory cytokines, reactive nitrogen species, and reactive oxygen species, characterizes this response [4]. In addition, lymphocytes are highly activated under chronic inflammatory conditions [5] the histology of inflamed tissue begins to resemble that of peripheral lymphoid organs, which can be referred to as lymphoid neogenesis or formation of tertiary lymphoid tissues. Lymphocyte recruitment to inflamed tissues is also reminiscent of lymphocyte homing to peripheral

^{*}Corresponding Author

Patricia Langjahr, Departmento de Biotecnología, Facultad de Ciencias Químicas, Universidad Nacional de Asunción, San Lorenzo, Paraguay. E-mail: plangjahr @ qui.una.py

^{© 2024} Christa Burgos *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (https://creativecommons.org/licenses/by/4.0/).

lymphoid organs. In the latter, under physiological conditions, homing receptors expressed on lymphocytes adhere to vascular addressin expressed on high endothelial venules (HEVs. Thus, overactivation of innate and adaptive responses can lead to chronic inflammation and autoimmune diseases.

Currently used anti-inflammatory drugs are effective; however, they have some disadvantages, such as undesirable adverse effects and unresponsiveness to treatment in certain patients [6]. Therefore, alternative treatment options are needed, and the study of plants with anti-inflammatory activity is relevant because of their potential pharmacological activity.

Baccharis (Asteraceae) is a widespread genus in the Americas and some species are commonly named "carqueja" [7]. *Baccharis* species are used in South American traditional medicine to treat gastrointestinal and liver disorders, infections, fever, and inflammation [8].

The *Baccharis* genus has been associated with numerous biological activities, including antioxidant, antibacterial, and antifungal properties [7,9–11]. Furthermore, previous studies have recognized potential anti-inflammatory effects [12–16]; however, its impact on the immune system has not been extensively analyzed.

Baccharitrimera is one of the most studied species within the genus *Baccharis*. The anti-inflammatory activity of *B. trimera* has been demonstrated in murine models of inflammation [12–14,17]. However, studies of its anti-inflammatory activity in human cells are limited.

Baccharis. punctulata has rarely been studied. It has been reported that *B. punctulata* exhibits anti-inflammatory properties in a murine model of 12-O-tetradecanoyl-phorbol-13-acetate-induced inflammation [18]. In addition, *B. punctulata* has been shown to decrease lymphocyte proliferation [19].

Therefore, in the present study, we analyzed the immunomodulatory activity of two species of *the Baccharis* genus, *B. punctulata* and *B. trimera*, in *in vitro* models of inflammation using monocytes and splenocytes, encompassing the assessment of both innate and adaptive immune responses.

MATERIALS AND METHODS

Plant material and sample preparation

Baccharis punctulata DC and B. trimera (Less.) DC were collected from Nemby, Paraguay, and San Lorenzo, Paraguay, respectively. Taxonomical evaluations were performed by botanists and voucher specimens (B. trimera: Voucher N° 4088 and B. punctulata: Voucher N° 4302) were deposited in the Herbarium-FCQ of the Department of Botany, Facultad de Ciencias Químicas, Universidad Nacional de Asunción (UNA). The aerial parts of the collected plants were used for extract preparation. The plant material was air-dried and ground into a fine powder using a cutting mill. To obtain the extracts, 200 g of the powder was suspended in HPLC-grade methanol (1.5 l) and sonicated three times for 30 minutes (inhouse method). The extraction procedure was repeated thrice, after which the extracts were filtered. A rotary evaporator (RVO 400 SD; Boeco, Germany) was used to remove the solvent. The yield was calculated considering the initial powdered dry material as the starting point (B. trimera: 17.1 g, 5.04%, and *B. punctulata*: 24.12 g, 6.96%). Plant methanol extracts were chemically characterized as previously described by our group [19]. The extracts were resuspended in dimethyl sulfoxide (DMSO; Sigma, USA) at a concentration of 10 or 100 mg/ml for further experiments.

Cells and cell culture

In this study, THP-1 (human acute monocytic leukemia) cells and splenocytes isolated from murine spleens were used. These cellular models were used to evaluate the immunomodulatory activity of the extracts on monocytes and lymphocytes, which are the primary cells involved in the innate and adaptive immune responses, respectively. To investigate the effect of *B. punctulata* and *B. trimera* extracts on innate immunity, their effects on human monocytic cells were determined using an *in vitro* model of lipopolysaccharide (LPS)-induced inflammation.

THP-1 cells were cultured at 37°C in 5% CO₂ in Roswell Park Memorial Institute (RPMI) medium (Gibco, USA), 10% fetal bovine serum (Gibco, Brazil), 100 U/ml penicillin, and 0.1 mg/ml streptomycin (Gibco, USA). Splenocyte isolation was performed as described in Coligan et al. [20] with minor modifications. Briefly, the spleen of 6-8-week-old healthy female Swiss albino mice were surgically removed in a sterile manner (under a laminar flow cabinet). The spleens were then gently pressed and subsequently subjected to a 70 µm cell strain. After that, the cells were incubated in red blood cell lysis using an ammonium chloride-potassium solution, washed with 1× phosphate buffer saline, and cultured in RPMI medium (Gibco, USA), 10% fetal bovine serum (Gibco, Brazil), 100 U/ ml penicillin, and 0.1 mg/ml streptomycin (Gibco, USA). Cell viability was assessed by trypan blue staining. The study was approved by the Ethics Research Committee of Facultad de Ciencias Químicas, UNA (N° 798/21).

Measurement of cell viability

THP-1 cells were seeded in 96-well plates (Corning Costar®, USA) at a density of 2×10^5 cells/well. The cells were treated with 10, 25, 50, 100, or 200 µg/ml of the extract for 24 hours. Cell viability was measured using an MTT colorimetric assay. Splenocytes, isolated as previously described, were cultured at a density of 5×10^6 cells/ml and 200 µl/well in a 96-well plate in RPMI medium in the presence of increasing extract concentrations. Resazurin was added 48 hours later and incubated for 24 hours at 37°C and 5% CO₂. After this time, cell viability was determined by absorbance measurements at 570 and 600 nm on a MultiskanTM GO (Thermo Scientific, Waltham, USA). Cells cultured in medium alone or 0.1% DMSO (vehicle control) were used as controls.

Determination of soluble CD14 (sCD14) levels

LPS induces monocyte activation and subsequent production of pro-inflammatory mediators such as cytokines and sCD14 [21–23] Cells (2×10^5 cells/well) were seeded in 96-well plates and pretreated for 15 minutes with the methanol extracts, followed by 1 µg/ml LPS (*E. coli* 0111:B4, Sigma, USA) for 24 hours. Cells treated with DMSO and cells pretreated with DMSO and stimulated with LPS were used as the negative and positive controls, respectively. sCD14 levels in the culture medium were assessed using an enzyme-linked immunosorbent assay (ELISA) assay (Quantikine ELISA, R&D Systems, USA) following the manufacturer's instructions. The optical density of each well was determined at 450 nm using a microplate reader (Multiscan GO; Thermo Fisher Scientific Waltham, USA).

Reverse transcription-polymerase chain reaction

THP-1 cells at a density of 1×10^6 cells/ml were treated with the methanolic extracts (15 minutes), followed by stimulation with LPS (1 µg/ml) for 4 hours as reported previously [24]. RNA extraction was performed using TRIzol reagent (Thermo Fisher Scientific, USA) and subsequently processed with DNase RQ1 (Promega, USA) at 37°C for 30 minutes. cDNA synthesis was performed with Moloney Murine Leukemia Virus Reverse Transcriptase (Promega®, USA), 100 µM Oligo dT, and dNTPs. cDNA was amplified (StepOnePlus real-time PCR system, Thermo Fisher Scientific, USA), using a SsoAdvanced Universal SYBR Green Supermix (BioRad, USA), and primers (Table 1).

A negative PCR control (with no template) was included. Relative mRNA expression was determined by the $2^{-\Delta\Delta Ct}$ method [29], using RPL13A as a normalizing gene.

Proliferation assay

Splenocytes (5 \times 10⁶ cells/ml) were seeded in RPMI medium and stimulated with 20 µg/ml concanavalin A (ConA) in the presence of extracts. After 48 hours of treatment, resazurin (AlamarBlue[®]) was added, and the absorbance at 570 and 600 nm was measured 24 hours later on MultiskanTM GO (Thermo Scientific, Waltham, USA). The proliferation index was calculated using the following equation:

$$IR = \frac{(\varepsilon_{OX})\lambda_2 A \lambda_1 - (\varepsilon_{OX})\lambda_1 A \lambda_2 \text{ extract}}{(\varepsilon_{OX})\lambda_2 A^{\circ} \lambda_1 - (\varepsilon_{OX})\lambda_1 A^{\circ} \lambda_2 DMSC}$$

Eox = molar extinction coefficient of oxidized resazurin; $\lambda_1 = 570$ nm; $\lambda_2 = 600$ nm.

Measurement of NO production

Griess assay [30] was performed using the culture supernatant of splenocytes treated with or without the extract for 15 minutes, followed by stimulation with LPS (10 μ g/ml) for 18 hours. The cell culture medium and Griess solution (0.1% N-1-naphthyl ethylenediamine dihydrochloride and 1% sulfanilamide in 5% phosphoric acid) were mixed in a 1:1 ratio and incubated for 15 minutes. The absorbance was measured at 545 nm (Multiscan GO, Thermo Fisher Scientific, USA). A standard curve was prepared by dissolving increasing concentrations of sodium nitrite (Cicarelli, Argentina) in the RPMI culture medium. The NO concentration was determined by extrapolation to the standard curve.

Statistical analyses

The results are expressed as mean \pm standard deviation and statistical significance was determined by one-way analysis of variance (ANOVA) and Dunnett's post-test (p < 0.05) using GraphPad Prism 8.01 (GraphPad Software Inc., USA).

RESULTS AND DISCUSSION

Cytotoxicity on THP-1 monocytic cells by *B. punctulata* and *B. trimera* extracts

First, the effect of the extracts on cell viability was determined by treating THP-1 cells with different concentrations of *B. punctulata* or *B. trimera* extracts. The species had no cytotoxic effects at 10, 25, or 50 μ g/ml (Fig. 1A and B) compared with cells treated only with the vehicle (DMSO) used to dissolve the plant extract. It should be noted that unstimulated cells (with culture medium only) showed no difference in viability from cells treated with the vehicle (DMSO). Based on these results, concentrations of up to 50 μ g/ml were selected for subsequent assays.

Effects of *B. punctulata* and *B. trimera* extracts on LPS-induced soluble CD14 and cytokines production in monocytic cells

THP-1 cells treated with the inflammatory stimulus LPS showed a significant increase in sCD14 production compared to unstimulated cells (Fig. 2). This indicated that the inflammation model functioned properly.

In addition, THP-1 cells were pretreated with vegetal extracts before stimulation with LPS. The levels of sCD14 in

Table 1. List of primers used for qPCR.

Gene	Primer direction	Primer sequence (5'-3')	Reference
RPL13A	Forward	CATCGTGGCTAAACAGGTACTG	[25]
	Reverse	CGCACGACCTTGAGGGCAGC	
IL-6	Forward	TCTCCACAAGCGCCTTCG	[26]
	Reverse	CTCAGGGCTGAGATGCCG	
MCP-1	Forward	GATCTCAGTGCAGAGGCTCG	[27]
	Reverse	TGCTTGTCCAGGTGGTCCAT	
IL-1β	Forward	TGATGGCTTATTACAGTGGCAATG	[28]
	Reverse	GTAGTGGTGGTCGGAGATTCG	

RPL13A: ribosomal protein L13a; IL-6: interleukin 6; MCP-1: monocyte chemoattractant protein-1;

IL-1β: interleukin 1 beta.



Figure 1. Effects of *B. punctulata* and *B. trimera* on cell viability. THP-1 cells were treated with *B. punctulata* (A) or *B. trimera* (B) extract for 24 hours. Cell viability was determined using an MTT assay. ANOVA: *p < 0.05, **p < 0.01, ***p < 0.001 versus vehicle (DMSO) control.



Figure 2. Inhibitory effects of *B. punctulata* and *B. trimera* extracts on the production of sCD14 in LPS-stimulated human monocytes. THP-1 cells were treated with *B. punctulata* (A) or *B. trimera* (B) extract in the presence of LPS (1 µg/ml) for 24 hours. ANOVA, $^{\#\#}p < 0.001$ versus untreated control; ***p < 0.001 versus LPS-treated cells.

the culture supernatant were subsequently measured, which revealed a decrease in sCD14 production in cells that were pretreated with the extracts and stimulated with LPS, compared to cells that were only treated with LPS. Therefore, treatment with *B. punctulata* or *B. trimera* markedly decreased LPSinduced sCD14 production (Fig. 2A and B), suggesting an immunomodulatory effect of these extracts on human monocytic cells. These results indicate a decrease in monocyte activation in the presence of the extracts compared to cells stimulated with LPS alone and without treatment with plant extracts. This suggests that the species modulates innate immunity.

LPS acts as a stimulus that triggers the production of inflammatory cytokines and chemokines in monocytes/ macrophages [22]. THP-1 cells were pretreated with B. punctulata or B. trimera extracts and subsequently stimulated with LPS. The cells were then collected, and RNA extraction and RT-qPCR were performed to evaluate IL-1, IL-6, and MCP-1 mRNA levels. Baccharipunctulata significantly reduced LPS-induced IL-6 and MCP-1 mRNA levels compared to those in cells stimulated with LPS (Fig. 3), indicating its antiinflammatory effects. Similarly, the B. trimera extract reduced LPS-induced IL-1ß and IL-6 mRNA levels (Fig. 4). Together, these results suggest that the extracts lead to a reduction in innate immune response. These findings are in line with the results observed in another species, B. dracunculifolia, which decreased IL-1 β , IL-6, and IL-10 levels in murine macrophages [15].

It should be noted that there were some differences in the effects of the species on cytokine expression: *B. punctulata* reduced IL-6 and MCP-1 mRNA levels, and *B. trimera* reduced IL-1 β and IL-6 mRNA levels, compared to those in cells stimulated with LPS alone. One explanation for this phenomenon is that different compounds were identified in these extracts in a previous study conducted by our group [19]. In our previous study, we showed that the methanol extract of *B. trimera* contained apigenin and genkwanin [19]. The flavonoid apigenin has been shown to decrease IL-6 and Il-1 β [31]. In addition, genkwanin has anti-inflammatory



Figure 3. Inhibition of cytokine production by the *B. punctulata* extract in LPS-stimulated human monocytes. THP-1 cells were treated with *B. punctulata* extract in the presence of LPS (1 µg/ml) for 4 hours. mRNA levels were analyzed by RT- RT-qPCR and the $2^{-\Delta \Delta Ct}$ method (each figure depicts fold change). ANOVA, "p < 0.05, "#p < 0.01, "##p < 0.01 versus untreated control; *p < 0.05, versus LPS-treated cells.



Figure 4. Inhibition of cytokine production by the *B. trimera* extract in LPSstimulated human monocytes. THP-1 cells were treated with *B. trimera* extract in the presence of LPS (1 µg/ml) for 4 hours. mRNA levels were analyzed by RT- RT-qPCR and the 2- $\Delta\Delta$ Ct method (each figure depicts fold change). ANOVA, "p < 0.05, ""p < 0.01, ""p < 0.05, ""p < 0.05, "**p < 0.001 versus untreated control; "p < 0.05, ***p < 0.001 versus LPS-treated cells.

activity and decreases IL-6 levels [32]. These compounds may be responsible for the action of the extract on the production of pro-inflammatory cytokines. In the *B. punctulata* methanol extract, trans-phytol and di-O-caffeoylquinic acid were previously identified by our group [19]. Di-O-caffeoylquinic acid and phytol have been shown to inhibit the production of the pro-inflammatory cytokines IL-6 and TNF- α [33,34]. Furthermore, di-O-caffeoylquinic acid has been shown to suppress the expression of inflammatory mediators such as nitric oxide(NO), prostaglandin E2, NO synthase, and cyclooxygenase-2 and to inhibit carrageenan-induced edema in vivo [33]. Therefore, these compounds could be responsible for the effects of the *B. punctulata* extract. These compounds have been documented to decrease the activation of the transcription factor NF- κ B, which stimulates the expression of pro-inflammatory cytokines and other inflammatory mediators [33,35]. Consequently, it is hypothesized that the mechanism of action of B. punctulata extract involves inhibition of NFκB activation.

Cytotoxicity on splenocytes and effects of *B. punctulata* and *B. trimera* extracts on cell proliferation

To gain a more comprehensive understanding of the effects of the extracts on immune cells, we evaluated the



Figure 5. Inhibitory effects of *B. punctulata* and *B. trimera* extracts on the proliferation of ConA-stimulated murine splenocytes. A-B. Effects of the extracts on cell viability. Splenocytes were treated with *B. punctulata* (A) or *B. trimera* (B) extract for 72 hours. ANOVA, **p < 0.01, ***p < 0.001 versus vehicle (DMSO) control. C-D. Splenocytes were treated with *B. punctulata* (A) or *B. trimera* (B) extract in the presence of ConA (20 µg/mL) for 72 hours. Splenocyte proliferation was determined using the Alamar Blue assay. ANOVA, **p < 0.001 versus DMSO control; ***p < 0.001 versus ConA-treated cells.

activities of *B. punctulata* and *B. trimera* in murine splenocytes. The extracts had no cytotoxic effects on splenocytes at concentrations of up to $50 \text{ }\mu\text{g/ml}$ (Fig. 5A and B).

ConA is a promoter of T lymphocyte division [36]. Therefore, ConA-induced splenocyte proliferation was determined as a marker of adaptive immunity [37]. Both species exhibited a decrease in the proliferation of cells induced by ConA (Fig. 5C and D), which further corroborated the antiinflammatory activity of these extracts.

In accordance with the present results, previous studies have shown that *B. trimera* inhibits phytohaemagglutinininduced human lymphocyte proliferation [14,19]. In addition, *B. punctulata* and *B. trimera* have been associated with a reduced production of IFN- γ , a cytokine produced by lymphocytes [19]. Our results and previous data suggest that *B. punctulata* and *B. trimera* affect T-lymphocyte proliferation.

Determination of *B. punctulata* and *B. trimera* effect on NO production

The effects of LPS-induced NO production were determined as representative indicators of the inflammatory response. *Baccharipunctulata* and *B. trimera* extracts decreased LPS-induced NO production (Fig. 6). These results indicate that both extracts decreased the inflammatory response pathway in murine splenocytes.



Figure 6. Effect of *B. punctulata* and *B. trimera* extracts on the production of nitric oxide (NO) in LPS-stimulated splenocytes. Cells were treated with *B. punctulata* or *B. trimera* extracts (50 µg/ml) in the presence of LPS (10 µg/ml) for 18 hours. NO production was determined by the Griess method. Data are expressed as the mean \pm SD. ANOVA. ^{##}p < 0.01 significance versus untreated control; *p < 0.05, versus LPS-treated cells.

NO, produced by immune cells, is an important mediator of the immune response. However, uncontrolled NO release can damage or destroy the host tissue. In the present study, *B. punctulata* and *B. trimera* reduced the LPS-induced NO production. Similarly, other *Baccharis* species (*B. obtusifolia, B. latifolia, B. pentlandii,* and *B. subulate*) have been shown to inhibit NO production [38]. Inhibition of iNOS activity, responsible for NO production, has been suggested to be beneficial for the treatment of pathologies involving inflammation [39]. Thus, *B. punctulata* and *B. trimera* present this potential action as immunomodulators of inflammatory response.

CONCLUSION

Evaluation of the activity of plant extracts on immune cells is a necessary step to recognize their potential in the management of inflammatory disorders such as autoimmunity, chronic infections, and allergies. In this study, we found that *B. punctulata* and *B. trimera* have anti-inflammatory activities, acting on both innate and adaptive immune responses. Our findings are significant because they show how these species globally modulate the inflammatory response.

In conclusion, the extracts tested in this study exhibited significant anti-inflammatory activity, acting on the innate and adaptive immune responses. Therefore, these species have important potential for the anti-inflammatory modulation of the inflammatory response.

AUTHOR CONTRIBUTIONS

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work. All the authors are eligible to be an author as per the International Committee of Medical Journal Editors (ICMJE) requirements/guidelines.

FINANCIAL SUPPORT

There is no funding to report.

CONFLICTS OF INTEREST

The authors report no financial or any other conflicts of interest in this work.

ETHICAL APPROVALS

The study protocol was approved by the Ethics Research Committee of Facultad de Ciencias Químicas, with approval number UNA (N° 798/21).

DATA AVAILABILITY

All data generated and analyzed are included in this research article.

USE OF ARTIFICIAL INTELLIGENCE (AI)-ASSISTED TECHNOLOGY

The authors declares that they have not used artificial intelligence (AI)-tools for writing and editing of the manuscript, and no images were manipulated using AI.

PUBLISHER'S NOTE

All claims expressed in this article are solely those of the authors and do not necessarily represent those of the publisher, the editors and the reviewers. This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

REFERENCES

- Ghasemian M, Owlia S, Owlia MB. Review of anti-inflammatory herbal medicines. Adv Pharmacol Sci. 2016;2016:9130979. doi: https://doi.org/10.1155/2016/9130979
- Kujawska M, Schmeda-Hirschmann G. The use of medicinal plants by Paraguayan migrants in the Atlantic Forest of Misiones, Argentina, is based on Guaraní tradition, colonial and current plant knowledge. J Ethnopharmacol. 2022;283:114702. doi: https://doi.org/10.1016/j. jep.2021.114702
- Medzhitov R. The spectrum of inflammatory responses. Science. 2021;374(6571):1070–5. doi: https://doi.org/10.1126/science. abi5200
- Kapellos TS, Bonaguro L, Gemünd I, Reusch N, Saglam A, Hinkley ER, *et al.* Human monocyte subsets and phenotypes in major chronic inflammatory diseases. Front Immunol. 2019;30(10):1–13. doi: https://doi.org/10.3389/fimmu.2019.02035
- Sakai Y, Kobayashi M. Lymphocyte "homing" and chronic inflammation. Pathol Int. 2015;65(7):344–54. doi: https://doi. org/10.1111/pin.12294
- Rice JB, White AG, Scarpati LM, Wan G, Nelson WW. Long-term systemic corticosteroid exposure: a systematic literature review. Clin Ther [Internet]. 2017;39(11):2216–29. doi: http://dx.doi. org/10.1016/j.clinthera.2017.09.011
- Ramos Campos F, Bressan J, Godoy Jasinski VC, Zuccolotto T, Da Silva LE, Bonancio Cerqueira L. *Baccharis* (Asteraceae): chemical constituents and biological activities. Chem Biodivers. 2016;13(1):1– 17. doi: https://doi.org/10.1002/cbdv.201400363

- Abad MJ, Bermejo P. *Baccharis* (Compositae): a review update. Arkivoc. 2006 Sep 5;2007(7):76–96. doi: http://dx.doi.org/10.3998/ ark.5550190.0008.709
- Pádua B da C, Silva LD, Rossoni JV, Humberto JL, Chaves MM, Silva ME, *et al.* Antioxidant properties of *Baccharis trimera* in the neutrophils of Fisher rats. J Ethnopharmacol. 2010;129(3):381–6. doi: https://doi.org/10.1016/j.jep.2010.04.018
- Gabaglio S, Alvarenga N, Cantero-González G, Degen R, Ferro EA, Langjahr P, *et al.* A quantitative PCR assay for antiviral activity screening of medicinal plants against Herpes simplex 1. Nat Prod Res. 2021;35(17):2926–30. doi: https://doi.org/10.1080/14786419.2 019.1675064
- Carrizo SL, Zampini IC, Sayago JE, Simirgiotis MJ, Bórquez J, Cuello AS, *et al.* Antifungal activity of phytotherapeutic preparation of *Baccharis* species from argentine Puna against clinically relevant fungi. J Ethnopharmacol. 2020;251:112553. doi: https://doi. org/10.1016/j.jep.2020.112553
- Gené RM, Cartañá C, Adzet T, Marín E, Parella T, Cañigueral S. Anti-inflammatory and analgesic activity of *Baccharis trimera*: identification of its active constituents. Planta Med. 1996;62(3):232– 5. doi: https://doi.org/10.1055/s-2006-957866
- Nogueira NPA, Reis PA, Laranja GAT, Pinto AC, Aiub CAF, Felzenszwalb I, *et al. In vitro* and *in vivo* toxicological evaluation of extract and fractions from *Baccharis trimera* with anti-inflammatory activity. J Ethnopharmacol. 2011;138(2):513–22. doi: https://doi. org/10.1016/j.jep.2011.09.051
- Paul EL, Lunardelli A, Caberlon E, De Oliveira CB, Santos RCV, Biolchi V, et al. Anti-inflammatory and immunomodulatory effects of *Baccharis trimera* aqueous extract on induced pleurisy in rats and lymphoproliferation *in vitro*. Inflammation. 2009;32(6):419–25. doi: https://doi.org/10.1007/s10753-009-9151-1
- Bachiega TF, de Sousa JPB, Bastos JK, Sforcin JM. Immunomodulatory/anti-inflammatory effects of *Baccharis dracunculifolia* leaves. Nat Prod Res. 2013 Sep;27(18):1646–50. doi: https://doi.org/10.1080/14786419.2012.742078
- Burgos C, Alfonso L, Ferro E, Langjahr P. Immunomodulatory activity of species of the genus *Baccharis*. Rev Paraguaya Reumatol. 2022;8(1):45–50. doi: 10.18004/rpr/2022.08.01.45
- de Oliveira CB, Comunello LN, Lunardelli A, Amaral RH, Pires MGS, da Silva GL, *et al.* Phenolic enriched extract of *Baccharis trimera* presents anti-inflammatory and antioxidant activities. Molecules. 2012;17(1):1113–23. doi: https://doi.org/10.3390/ molecules17011113
- Ascari J, de Oliveira MS, Nunes DS, Granato D, Scharf DR, Simionatto E, *et al.* Chemical composition, antioxidant and antiinflammatory activities of the essential oils from male and female specimens of *Baccharis punctulata* (Asteraceae). J Ethnopharmacol. 2019;234:1–7. doi: https://doi.org/10.1016/j.jep.2019.01.005
- Burgos C, Alvarenga N, Heiderich H, Florentín-pavía M, Sotelo PH, Carpinelli MM, *et al.* Immunomodulatory effects of three species of *Baccharis* on human peripheral blood mononuclear cells. Trop J Nat Prod. 2021;1055–9. doi: https://doi.org/10.26538/tjnpr/v5i6.12
- Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, Strober W. Current protocols in immunology. Hoboken, NJ: John Wiley and Sons; 1991. Available from: https://www.wiley.com/en-au/ Current+Protocols+in+Immunology+-p-9780471522768
- 21. Ma WT, Gao F, Gu K, Chen DK. The role of monocytes and macrophages in autoimmune diseases: a comprehensive review. Front Immunol. 2019;10:1–24. doi: https://doi.org/10.3389/fimmu.2019.01140
- Ngkelo A, Meja K, Yeadon M, Adcock I, Kirkham PA. LPS induced inflammatory responses in human peripheral blood mononuclear cells is mediated through NOX4 and G iα dependent PI-3kinase signalling. J Inflamm. 2012;9:2–8. doi: https://doi.org/10.1186/1476-9255-9-1
- Zanoni I, Granucci F. Role of CD14 in host protection against infections and in metabolism regulation. Front Cell Infect Microbiol. 2013;3(32):1–6. doi: https://doi.org/10.3389/fcimb.2013.00032

- Chanput W, Mes J, Vreeburg RAM, Savelkoul HFJ, Wichers HJ. Transcription profiles of LPS-stimulated THP-1 monocytes and macrophages: A tool to study inflammation modulating effects of food-derived compounds. Food Funct. 2010;1(3):254–61. doi: https://doi.org/10.1039/c0fo00113a
- Yamada S, Kotake Y, Demizu Y, Kurihara M, Sekino Y, Kanda Y. NAD-dependent isocitrate dehydrogenase as a novel target of tributyltin in human embryonic carcinoma cells. Sci Rep. 2014;4:5952. doi: https://doi.org/10.1038/srep05952
- Wolf K, Schulz C, Riegger GAJ, Pfeifer M. Tumour necrosis factor-α induced CD70 and interleukin-7R mRNA expression in BEAS-2B cells. Eur Respir J. 2002;(2):369–75. doi: https://doi.org/10.1183/09 031936.02.00303602
- Wang H, Yi T, Zheng Y, He S. Induction of monocyte chemoattractant protein-1 release from A549 cells by agonists of protease-activated receptor-1 and -2. Eur J Cell Biol. 2007;86(4):233–42. doi: https:// doi.org/10.1016/j.ejcb.2006.12.003
- Ando Y, Kuroda A, Kusama K, Matsutani T, Matsuda A, Tamura K. Impact of serine protease inhibitor alpha1-antitrypsin on expression of endoplasmic reticulum stress-induced proinflammatory factors in adipocytes. Biochem Biophys Reports. 2021;26:100967. doi: https:// doi.org/10.1016/j.bbrep.2021.100967
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. Methods. 2001;(4):402-8. doi: https://doi.org/10.1006/meth.2001.1262
- Stich RW, Shoda LKM, Dreewes M, Adler B, Jungi TW, Brown WC. Stimulation of nitric oxide production in Macrophages by *Babesia bovis*. Infect Immun. 1998;66 (9):4130–6. doi: https://doi. org/10.1128/IAI.66.9.4130-4136.1998
- Zhang X, Wang G, Gurley EC, Zhou H. Flavonoid apigenin inhibits lipopolysaccharide-induced inflammatory response through multiple mechanisms in Macrophages. PLoS One. 2014;9(9):e107072. doi: https://doi.org/10.1371/journal.pone.0107072
- Gao Y, Liu F, Fang L, Cai R, Zong C, Qi Y. Genkwanin inhibits proinflammatory mediators mainly through the regulation of miR-101/MKP-1/MAPK pathway in LPS-activated macrophages. PLoS One. 2014;9(5):e96741. doi: https://doi.org/10.1371/journal. pone.0096741

- Jang G, Lee S, Hong J, Park B, Kim D, Kim C. Anti-inflammatory effect of 4,5-dicaffeoylquinic acid on raw264.7 cells and a rat model of inflammation. Nutrients. 2021;13(10):1–10. doi: https://doi. org/10.3390/nu13103537
- Zai JA, Khan MR, Mughal ZUN, Khan FS, Soomro N. Phytol anti-oxidative and anti-inflammatory effects in hydrogen peroxide challenged human PMBCs involves NF κ B pathway. Int J Emerg Technol. 2021;12(1):296–303.
- 35. Carvalho AMS, Heimfarth L, Pereira EWM, Oliveira FS, Menezes IRA, Coutinho HDM, *et al.* Phytol, a chlorophyll component, produces antihyperalgesic, anti-inflammatory, and antiarthritic effects: possible NFκB pathway involvement and reduced levels of the proinflammatory cytokines TNF-α and IL-6. J Nat Prod. 2020;83(4):1107–17. doi: https://doi.org/10.1021/acs. jnatprod.9b01116
- Dwyer JM, Johnson C. The use of concanavalin A to study the immunoregulation of human T cells. Clin Exp Immunol. 1981;46(2):237–49.
- Bonilla FA, Oettgen HC. Adaptive immunity. J Allergy Clin Immunol. 2010;125(2):S33–40. doi: https://doi.org/10.1016/j. jaci.2009.09.017
- Abad MJ, Bessa AL, Ballarin B, Arag O, Gonzales E, Bermejo P. Anti-inflammatory activity of four Bolivian *Baccharis* species (Compositae). J Ethnopharmacol. 2006;103:338–44. doi: https://doi. org/10.1016/j.jep.2005.08.024
- Wong VC, Lerner E. Nitric oxide inhibition strategies. Futur Sci OA. 2015;1(1):FSO35. doi: https://doi.org/10.4155/fso.15.35

How to cite this article:

Burgos C, Alvarenga N, Sotelo PH, Langjahr P. Suppression of inflammation by *Baccharis punctulata* and *Baccharis trimera* through modulation of innate and adaptive immune responses. J Appl Pharm Sci. 2024;14(08):078–084.