



Journal of Applied Pharmaceutical Science

Available online at www.japsonline.com

ISSN: 2231-3354
Received on: 04-05-2012
Revised on: 12-05-2012
Accepted on: 18-05-2012
DOI: 10.7324/JAPS.2012.2538

The effect of ginger and garlic on the microbial load and shelf life of Kunun-zaki

Edward, K.C. and Ohaegbu, C.G.

ABSTRACT

Kunun-zaki was produced using ginger and garlic and stored under ambient conditions for 10 days. The microbiological load and the shelf life of the drink were investigated. Diverse microbial genera: *Lactobacillus*, *Bacillus*, *Staphylococcus*, *Aspergillus*, *Penicillium*, *Fusarium* and *Saccharomyces* were isolated from samples. The effect of ginger and garlic separately were compared to the combined effect of ginger and garlic in reducing the microbial population. Of all the treatments, garlic (2g in 200mls or 0.01% w/v) was most effective in reducing the microbial populations. In contrast, treatment with 1g of ginger was least effective in reducing the microbial populations. Shelf life based on sensory overall acceptability and microbial quality of the samples varied with treatments but combination treatment with 2g in 200mls(0.01% w/v) ginger and garlic extended the shelf life by approximately four (4) days whereas other treated samples showed marginal enhanced shelf life of 2days. However, untreated control sample exhibited remarkably high microbial load and was virtually unacceptable after 24h of production. The result shows the potential of the combination treatment of ginger and garlic as antimicrobials and in extending the shelf life of Kunun-zaki.

Keywords: Kunun-zaki, ginger, garlic, shelf life, microbial load.

INTRODUCTION

Kunun-zaki, a non alcoholic beverage, very popular, especially in Northern Nigeria is known for its social, religious and therapeutic values. It is taken as a refreshing drink, appetizer, food compliment, thirst quencher, substitute for or to complement soft drinks and wines at social gatherings. (Oramili *et al*; 2003). The nutritional value of Kunun-zaki as documented by Ayo and Okaka (1998) and Sopade and Kassun (1992) justifies its use. It is found to be rich in carbohydrate, vitamins and minerals and contains protein and fat. Sweet potato, which is used as an additive in its preparation, adds to the vitamin content. Kunun-zaki is produced from fermented millet, sorghum, guinea corn and maize in decreasing order of preference and is normally flavoured with ginger. It is consumed in the active state of fermentation.

Edward, K.C. and Ohaegbu, C.G.
Department of Microbiology,
Michael Okpara University of
Agriculture, Umudike, Nigeria.

For Correspondence
Edward, K.C
Department of Microbiology,
Michael Okpara University of
Agriculture, Umudike, Nigeria
Phone: +234-703 3420 330

The fermentation process is normally dominated by the microbial genera; *Lactobacillus*, *Bacillus*, *Aspergillus* and *Saccharomyces*. (Efiuvwevwere and Akoma, 1997). The method of preparation is simple and cheap and does not require any elaborate equipment. (Agboola, 1987) The short shelf life of Kunun-zaki is a major problem of both the brewers and consumers of this drink. These deleterious changes are due primarily to the objectionable off-flavour (over souring) induced by microorganisms. The use of chemical preservatives for microbial control in beverages is desirable and continues to generate research interest worldwide. However, with most of the published information being on the use of inorganic chemical preservatives to preserve this product, the objective of this study is therefore based on the use of organic chemical preservatives (ginger and garlic) to enhance the shelf-life stability of the product while maintaining the characteristic organoleptic attributes.

MATERIALS AND METHODS

Sample collection

Millet grains (*Pennisetum typhoides*), sugar, dried pepper, ginger and garlic were obtained from Umuahia main market, Abia State, Nigeria.

Production

Kunun-zaki was produced using the method described by Efiuvwevwere and Akoma (1995). Millet grains (1000g) were cleaned and steeped in water (1.5L) for 24h at ambient temperature. After 24h, the water was decanted off and the grains washed with tap water before blending with 10g of dried pepper in two volume tap water. The slurry was sieved and the filtrate allowed to sediment for 3-5h at ambient temperature. The supernatant was discarded while the various treatments were added to the remaining milky and pasty sediment. Each of the different concentrations of the treated samples and the control were divided into two portions. One of these was gelled using two volume boiling water and then allowed to cool to between 45°C -50°C. The gelled portions of each of the different concentrations of the samples and control were mixed (1:1 by volume) with the ungelled portion. Each of the mixture was then diluted with tap water, allowed to ferment for 8h and then sweetened with granulated sugar. The samples were dispensed into sterile screw capped containers.

Storage

The samples were stored at ambient temperature (25-32°C) for 10 days.

Enumeration and isolation of microorganisms

This was carried out on Nutrient agar (NA), Sabouraud Dextrose agar (SDA), and Rogosa agar (oxoid) using pour plate method. The samples were serially diluted and 1ml of appropriate dilution was used to inoculate each of the agar plates in triplicates. The nutrient and rogosa agar plates were incubated at 37°C aerobically for 24-36h while the SDA plates were incubated at

room temperature for 48-72h. Colonies were counted using a colony counter (Gallenkamp). The mean of duplicate results were then recorded as the colony count. (Lateef *et al.*, 2004).

Identification of isolates

Discrete colonies were picked at random, purified and characterized based on standard microbiological cultural, morphological and biochemical characteristics. (Harrigan and McCance, 1976 and Sneath *et al.*, 1986). The biochemical tests include; catalase, oxidase, Voges-Proskauer, hydrogen sulphide production, nitrate reduction, citrate utilization and sugar fermentation.

For the fungi, the colonies were screened and identified based on the taxonomic schemes and descriptions by Fawole and Oso, (1998) and Mislivec *et al.*, (1992)

Sensory analysis

The sensory attributes were evaluated as described by Larmond (1977).

Physicochemical analysis

pH and titrable acidity were determined. The pH was determined using referenced pH meter (model 291MK2, PYE UNICAM, England) while the titrable acidity was determined using the method described by Speck (1984)

RESULTS AND DISCUSSION

The rapid deterioration in shelf life of traditionally produced Kunun-zaki and other African beverages is widely acknowledged and is of great concern (Odunfa, 1985; Dirar, 1993; Efiuvwevwere and Akoma, 1995). The occurrence of diverse microbial genera and the remarkable high microbial load in the untreated control sample (Tables 1,2 and 3) are major cause of accelerated spoilage commonly experienced by brewers and consumers of these products. It is evident from tables 1,2 and 3 that the treated samples have lower microbial load because the added spices (ginger and garlic) have antimicrobial effect and are capable of destroying pathogenic bacteria (Ayo *et al.*, 2003).

In general, ginger and garlic have comparable antimicrobial activities due to the presence of essential oils in them (Nakatani, 2000, Ilondu *et al.*, 2001). But the more appreciable inhibitory effect exhibited by garlic (Tables 1,2, and 3) may be attributed to the differences in their essential oil components. Garlic tends to exert a more pronounced cell membrane interface and disruption than ginger due to the action of Allistatin I and Allistatin II (diallyldisulphide oxide) contained in it which are not present in ginger and which affect the growth and respiration of microorganisms (Tynecka and Gos, 1973). Whereas the treatment with ginger and garlic separately resulted in reduction in the microbial load, the combination of the two (ginger and garlic) showed comparable efficacy when compared to that of treatment with ginger alone.

However, the magnitude of effectiveness of ginger and garlic as organic preservative on the microbial load of the sample

differed with the concentration. A decrease in the number of microorganisms was observed with the increase in the concentration of the preservatives. But the effects of ginger and garlic tended to decrease with storage time and this could be due to microbial degradation particularly in the presence of high lactobacilli load (De Boer, 1998).

Similar findings had been reported by Efiuvwevwe and Akoma, (1997) during Kunun-zaki preservation using inorganic preservatives. The drop in pH of all samples over the six days of storage (Table 4.) was in line with what was reported by Agarry *et al.*(2010). This may be as a result of greater microbial activities since this decrease was more observed in the untreated control samples having high microbial load. The high rate of change in pH with storage days could be due to the decomposition of fermentable substrates and sugars by microorganisms especially *Lactobacillus* species which ferment carbohydrates to produce energy and principally lactic acid.

The sensory analysis showed that the treated samples were more acceptable than the untreated control sample. The high rate of acceptability by the 10-member panel for the treated samples is attributed to the extra flavour added by the spices (Adeyemi and Umar, 1994). Combination treatment of ginger and garlic were more acceptable by the panelists with acceptability increasing with concentration and it extended the shelf life by about four (4) days. However, Kunun-zaki with garlic treatment

alone showed apparent beneficial antimicrobial effects, but consumer acceptability was lower, confirming the off flavour that occurred when the concentration was increased.

The bacteria isolated included *Staphylococcus aureus*, *Lactobacillus* and *Bacillus* species. The dominance of *Lactobacillus* species in the samples may be due to the acidic nature of the drink and the presence of ginger is known to encourage its growth. Also, this could be as a result of one or more substances such as organic acids, ethanol, hydrogen peroxide and bacteriocins produced by *Lactobacillus* species during growth which inhibit or destroy the growth of other competitive microorganisms and thus improving the safety and keeping quality of such food products. This dominance made Efiuvwevwe and Akoma (1997) to conclude that Kunun-zaki is a lactic acid bacteria fermented product. The presence of *Bacillus* species in the samples is not accidental as they occur always on food of low acid content like drinks where they produce organic acids. They may also enter by soil contamination.

Consequently, the isolation of *Staphylococcus aureus* may be attributed to processing and post processing contamination through processors and utensils. Fungi isolated were *Aspergillus*, *Penicillium*, *Fusarium* and *Saccharomyces* species. The presence of these fungi is associated with spoilage of beverages. *Saccharomyces* spp are known to convert starch and dextrin to glucose which is then fermented to ethanol and carbon dioxide.

Table. 1: Effect of ginger and garlic on the total viable count of Kunun-zaki during tropical ambient storage.

Storage (day)	K+GG 1g	K+GG 2g	K+GL 1g	K+GL 2g	K+GG+GL 0.5g each	K+GG+GL 1g each	Control
	10^4 cfu/ml						
0	4.9	4.7	4.3	3.8	4.6	4.2	5.3
2	6.5	5.5	5.4	4.5	5.7	5.0	7.2
4	2.8	25	26	22	27	24	ND
6	30	28	28	25	26	27	ND
8	28	26	25	23	26	25	ND
10	23	21	19	16	20	21	ND

KEY:

K = Kunun-zaki, GG = Ginger, GL = Garlic, Control (contains neither ginger nor garlic), ND = Not Determined

Table. 2: Effect of ginger and garlic on the Total fungal counts of Kunun-zaki.

Storage (day)	K+GG 1g	K+GG 2g	K+GL 1g	K+GL 2g	K+GG+GL 0.5g each	K+GG+GL 1g each	Control
	10^4 cfu/ml						
0	4.0	3.9	3.4	3.0	3.6	3.5	4.3
2	5.8	5.0	4.2	3.3	5.0	3.9	64
4	22	19	16	15	19	17	26
6	26	23	19	17	25	20	30
8	16	13	13	11	14	12	24
10	12	10	11	10	12	10	20

KEY:

KZ = Kunun-zaki, GG = Ginger, GL = Garlic, Control (contains neither ginger nor garlic), ND = Not Determined

Table. 3: Effect of ginger and garlic on the *Lactobacilli* count.

Storage (day)	K+GG 1g	K+GG 2g	K+GL 1g	K+GL 2g	K+GG+GL 0.5g each	K+GG+GL 1g each	Control
	10^4 cfu/ml						
0	4.3	4.1	3.6	3.2	3.9	3.8	4.6
2	6.0	5.2	4.5	3.5	5.3	4.0	70
4	28	26	25	18	27	22	30
6	30	29	29	22	29	26	ND
8	27	25	23	20	25	23	ND
10	21	17	17	13	19	16	ND

KEY:

KZ = Kunun-zaki, GG = Ginger, GL = Garlic, Control (contains neither ginger nor garlic), ND = Not Determined

Table 4: Effect of ginger and garlic on the pH of Kunun-zaki.

Storage (day)	K+GG 1g	K+GG 2g	K+GL 1g	K+GL 2g	K+GG+GL 0.5g each	K+GG+GL 1g each	Control
0	4.65	4.69	4.74	4.82	4.70	4.76	4.62
2	4.05	4.09	4.14	4.20	4.10	4.16	3.56
4	3.01	3.04	3.10	3.18	3.06	3.12	2.20
6	2.25	2.29	2.33	2.38	2.29	2.34	1.20
8	2.20	2.22	2.25	2.30	2.23	2.27	ND
10	2.10	2.14	2.18	2.20	2.14	2.16	ND

KEY:

KZ = Kunun-zaki, GG = Ginger, GL = Garlic, Control (contains neither ginger nor garlic), ND = Not Determined

Table 5: Effect of ginger and garlic on overall sensory acceptability of Kunun-zaki.

Storage (day)	K+GG 1g	K+GG 2g	K+GL 1g	K+GL 2g	K+GG+GL 0.5g each	K+GG+GL 1g each	Control
0	2.44	2.48	2.50	2.46	2.30	2.10	2.25
2	3.20	3.10	3.00	2.84	2.68	2.60	4.62
4	5.80	5.48	5.26	4.40	3.25	3.06	6.50
6	6.08	6.00	5.89	5.70	5.52	5.40	7.00
8	7.22	7.08	6.88	6.40	6.15	6.10	ND
10	ND	ND	ND	7.50	7.20	7.02	ND

KEY:

KZ = Kunun-zaki, GG = Ginger, GL = Garlic, Control (contains neither ginger nor garlic), ND = Not Determined

CONCLUSION

From this study, the combined treatment of garlic and ginger, especially the 0.01(w/v) extended the shelf life of Kunun zaki up to about 4days. The shelf life extension means enhanced commercial potential of Kunun zaki. The combined treatment with ginger and garlic especially the 0.01 (w/v) is therefore recommended.

REFERENCES

- Adeyemi, T. Umar, S. Effect of manufacture on the quality characteristics of Kunun-zaki, a millet based beverage. *Nigerian Food Journal*. 1994; 12:34-40
- Agari, O.O., Nkama, I. Akoma, O. Production of Kunun zaki (A Nigerian fermented cereal beverage) using starter culture. *International Research Journal of Microbiology*. 2010; 1(2):018-025
- Agboola, S.D. Storage and preservation of Agricultural products in Nigeria, current status of methods. Seminar on integrated rural development in Kwara State, Ilorin A.A. Ajaani, O. Bello (Eds.). Directorate of Foods, Roads and Rural infrastructure. (1987) 10-15
- Ayo, J.A. and Okaka, J.C. Interaction effect of Candaba farinose extract and pH levels on some physicochemical properties of Kunun-zaki. Proceedings of the 22nd Annual NIFST Conference, Abeokuta, Nigeria. (1998) 31-33.
- Ayo, J.A., Ejiogu, T.N. Iheanacho. Effect of spices on the microbial and sensory quality of Kunun-zaki. Proceedings of the 27th Annual NIFST Conference. (2003) 233
- DeBoer, E. Introduction to food borne fungi (3rd ed). Central bureau Schimmelcultures, Baam. (1998) 268-273.
- Dirar, HA The buligenous fermented foods of the Sudan. A study in African foods and Nutrition, Wallingford, Cambridge University press. (1993) 34-36.
- Efiuwwewere, B. J. O., Akoma, O. The microbiology of kunun-zaki, a cereal beverage from Northern Nigeria during the fermentation (production) process. *World Journal of Microbiology and Biotechnology*. 1995; 1:491-493.
- Efiuwwewere, B, J, O., Akoma, O. Effects of chemical preservative and pasteurization on the microbial spoilage and shelf-life of kunun-zaki. *Journal of Food Safety*. 1997;17:203-213.

Fawole, MO, Oso, BA Laboratory Manual of Microbiology. Spectrum books Limited, Ibadan. (1998) 26-31

Harrigan, WF, McCance, ME. Laboratory Methods in Food and Dairy Microbiology. Academic press, London. (1976) 42.

Halzapfel, W. H. (1989) Industrialization of African indigenous beer brewing: In industrialization of Indigenous Fermentation Foods (K. H. Steinkraus, Ed.) 191-283. New York; Marcel Dekker Inc.

Ilondu, E. M., Ejechi, B. O., Souzey, J. A., Microbial stability of jam prepared from Velvet tamarind and preserved by combined process. *Nigerian Journal of Microbiology*. 2001; 5:93-96.

Larmond, E. Laboratory methods for Sensory Evaluation of Foods. Canada Department of Agriculture, Ottawa. Canada. (1977) Pp. 12-88.

Maji, A.A. Omale, J. and Okoli, E.C. Effects of chemical preservatives and pasteurization on the shelf life of Kunun zaki (sorghum and maize gruel) *European Journal of Food Research and review*. 2011; 1(2): 61-70

Mislivec, P.B., Beuchat, L.R., Cousin, M.A *Yeasts and molds* In: Compendium of methods for the Microbiological examination of foods. C. Vanderzant, and D.F Splittstoesser, (Eds.). American Public Health Association, Washington, D.C. (1992)

Nakatani, N. Phenolic antioxidants from herbs and spices *Journal of Biofactors*. 2000; 13: 141-146.

Odufa, S. A. (1985) African fermented foods In: *Microbiology of Fermented foods*. Vol.2. B.J.B. Wood (Ed.) pp 155-191. London: Elsevier Applied Science.

Oranusi, S. U., Umoh, V. J., Kwaga, J. K. P. Hazards and critical control points of kunun-zaki, a non-alcoholic beverage in Northern Nigeria. *Food Microbiology*. 2003; 20: 127-132.

Sneath, P. H. A., Nair, N. S., Sharpe, N. E., Holt, J. G. *Bergey's Manual of systemic Bacteriology*. Vol.2, Williams and Wilkins, Baltimore, M. D. 1986; pp23-100.

Sopade, P. A., Kassum, A. L. Rheological characterization of Nigerian liquid and semi-solid foods Kunun-zaki and kunun-gyada. *Nigerian Food Journal*. 1992; 10:23-33.

Speck, ME. Compendium of methods for the Microbiological Examination of foods, 2nd Ed. American public Health Association, Washington, D.C. (1984) 56-90.

Tynecka, Z., Gos, Z. The inhibition action of Garlic (*Allium Sativum*) on growth and respiration of some micro-organisms. *Acta Microbiology Polinica*. 1973; Series B 5:51-62.