

**Expression of polyprotein and 3D polymerase protein in Sf9 cells and immunogenicity against enterovirus A71B5 (Thailand strain)**

**Nipatha Issaro<sup>1\*</sup>, Aphisak Kongkaew<sup>2</sup>, Akanitt Jittmittraphap<sup>3</sup>, Pornsawan Leungwutiwong<sup>3</sup>, Wutigri Nimlamool<sup>4</sup>, Mingkwan Na Takuathung<sup>4</sup>**

<sup>1</sup>Department of Pharmacognosy and Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Burapha University, Chonburi, Thailand.

<sup>2</sup>Research Administration Section, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand.


<sup>3</sup>Department of Microbiology and Immunology, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

<sup>4</sup>Department of Pharmacology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand.


doi: <https://doi.org/10.7324/JAPS.2023.93192>

**Supplementary Material**

**Supplementary Figures:**



**Fig. S1.** SDS-PAGE analysis of P1-3D<sup>pol</sup> purification. Lane M: Protein marker; lane FT: Flow through; lane W1-W4: Washing Buffer at 10, 20, 50, and 100 mM imidazole, respectively.



**Fig. S2.** Western blot analysis of purified P1-3D<sup>pol</sup>. The result show that the peptides/proteins with molecular weights near 35-45 kDa and 60-70 kDa could be recognized by VP0 antibodies (Merck, Millipore MAB979), and might be EV71 proteins.