

Synthesis, *in silico*, and *in vitro* pharmacological evaluation of norbornenylpiperazine derivatives as potential ligands for nuclear hormone receptors

Karine Badalyan¹, Nelly Babayan^{2,3}, Elena Kalita^{2,3}, Narine Grigoryan², Natalya Sarkisyan², Ruzanna Grigoryan², Grigor Arakelov⁴, Alexan Shahkhatuni⁵, Hovhannes Attaryan¹, Davit Mkrtchyan¹, Hasmik Khachatryan¹, Lusine Khondkaryan²

¹Laboratory of Applied Chemistry, Scientific and Technological Center of Organic and Pharmaceutical Chemistry NAS RA, Yerevan, Republic of Armenia.

²Laboratory of Cell Technologies, Institute of Molecular Biology NAS RA, Yerevan, Republic of Armenia.

³Laboratory of Experimental Biology, CANDLE Synchrotron Research Institute, Yerevan, Republic of Armenia.


⁴Laboratory of Computational Modelling of Biological Processes, Institute of Molecular Biology NAS RA, Yerevan, Republic of Armenia.

⁵Molecule Structure Research Center, Scientific and Technological Center of Organic and Pharmaceutical Chemistry NAS RA, Yerevan, Republic of Armenia.


Doi: <http://dx.doi.org/10.7324/JAPS.2025.230239>

SUPPLEMENTARY MATERIALS


Figure S1. ¹HNMR spectrum of compounds 4a-4h.




4b




4c




4d




4e



4f



4g



4h






Figure S2. ¹³C NMR spectrum of compounds 4a-4h (a-h).


4a




4b




4c




4d




4e



4f



4g



4h






Figure S3. IR spectra of compounds 4a-4h.


4a




4b




4c




4d




4e



4f



4g



4h




Figure S4. Mass spectral analysis of compound 4b.




Figure S5. Mass spectral analysis of compound 4d.

